

AN EXAMPLE OF AN INDECOMPOSABLE MODULE WHICH IS NOT INJECTIVE

Dedicated to Professor Manabu Harada on his 60th birthday

SHOJI MORIMOTO and TAKEFUMI SHUDO

It is well-known that every torsionfree divisible module is injective over a commutative integral domain. First we show that this theorem can be improved by using the concept of weakly σ -injective modules over a left Ore domain. Also in [2], we show that for a non-singular module M if M has no nonzero injective submodule, then so does M^Λ for all index sets Λ . Finally, we give an example to show that the above proposition is false in general.

Throughout this note R is a ring with identity and modules are unitary left R -modules unless otherwise stated. We denote the category of modules by $R\text{-mod}$ and the injective hull of a module M by $E(M)$. As for terminologies and basic properties concerning torsion theories and preradicals, we refer to [3]. Let ρ be a preradical. We call it *stable* if $T(\rho)$ is closed under essential extensions. Also the left linear topology corresponding to a left exact preradical ρ is denoted by $\mathcal{L}(\rho)$. Now for two preradicals ρ and τ , we shall say that ρ is *larger than* τ if $\rho(M) \supseteq \tau(M)$ for all modules M .

We put $\mathcal{D} = \{r \in R \mid rs \neq 0 \text{ and } sr \neq 0 \text{ for all } s (\neq 0) \in R\}$. We call a ring R *left Ore* if for each $r \in R$ and $s \in \mathcal{D}$ there exist $r' \in R$ and $s' \in \mathcal{D}$ such that $s'r = r's \neq 0$. Also we put $\sigma(M) = \{x \in M \mid rx = 0 \text{ for some } r \in \mathcal{D}\}$. In general, $\sigma(M)$ is not a submodule of M .

However

Proposition 1. *If R is a left Ore domain, then σ is the Goldie torsion functor G .*

Proof. By [3, p. 138, Example 2], σ is a left exact radical. By assumption, every non-zero ideal of R is essential in R . Thus $\mathcal{L}(Z)$ is the set of non-zero ideal of R , where Z is the singular torsion functor. Thus $\sigma \leq Z$. Conversely let r be a non-zero element of R and let $s + Rr$ be in R/Rr . Then $s's = r'r$ for some r' and s' in R by assumption. Thus R/Rr is in $T(\sigma)$ and so Rr is in $\mathcal{L}(\sigma)$. Hence every non-zero ideal of R belongs to $\mathcal{L}(\sigma)$. Since σ is a radical, so is Z , namely, $Z = G$. Hence $\sigma = G$.

Definition. Let τ be a preradical. We call a module H (resp. *weakly*) τ -*injective* if for all exact sequences of modules $O \rightarrow A \rightarrow B \rightarrow C \rightarrow O$ with $C \in T(\tau)$ (resp. $B \in T(\tau)$), the functor $\text{Hom}_R(-, H)$ preserves the exactness.

Lemma 2 [1, Theorem 1. 11.]. *Let τ be a left exact preradical. Then the following conditions are equivalent :*

- (i) *Every τ -injective module is injective.*
- (ii) *$\bar{\tau}$ is larger than the Goldie torsion functor G , where $\bar{\tau}$ is the smallest radical larger than τ .*

By the above lemma, every σ -injective module is injective.

Theorem 3. *Let R be a left Ore domain. For a module M , the following conditions are equivalent :*

- (i) *M is divisible and weakly σ -injective.*
- (ii) *M is injective.*

Proof. (ii) \Rightarrow (i) is clear. (i) \Rightarrow (ii). By Proposition 1, $\sigma = G$. Thus it is sufficient to show that M is σ -injective by Lemma 1.2. We assume that $\sigma(E(M)/M) \neq O$. Then there exists $\bar{x} = x + M$ (x is in $E(M)$ and is not in M) such that $r\bar{x} = 0$ for some $r (\neq 0)$ in R . Since M is divisible and rx is in M , $rx = rm$ for some m in M , namely $r(x - m) = 0$. Thus $x - m$ is in $\sigma(E(M))$. Since M is weakly σ -injective, $\sigma(E(M)) = \sigma(M)$ and so $x - m$ is in M . Hence x is in M . This is a contradiction. Thus M is σ -injective.

Since σ is left exact, every σ -torsionfree module is weakly σ -injective. Thus we have the following famous result :

Corollary 4. *Let R be a commutative integral domain and M a torsionfree module. Then M is injective if and only if it is divisible.*

We call a ring R *left hereditary* if every left ideal of R is projective. From Theorem 3 and [3, Proposition 4.5], we have

Corollary 5. *Let R be a left Ore domain. Then the following conditions are equivalent :*

- (i) *Every divisible module is weakly σ -injective.*
- (ii) *Every divisible module is σ -injective.*
- (iii) *Every divisible module is injective.*
- (iv) *R is left hereditary.*

First we give an example of a module which is divisible but not injective.

Let \mathbf{Z} be the ring of integers, \mathbf{Q} the field of rational numbers and \mathbf{Z}_p the localization of \mathbf{Z} with respect to $p\mathbf{Z}$ for a prime number p .

Example 6. *Let R be the polynomial ring over \mathbf{Z}_p and K the quotient field of R . Then K/R is not injective.*

Proof. We put $I = pR + xR$. Let f be a map from I to K/R with $f(pa_0 + a_1x + \dots + a_nx^n) = (p/x)(a_0 + a_1x + \dots + a_nx^n) + ((1-p)/p)(a_1 + a_2x + \dots + a_nx^{n-1}) + R$, where $a_i (i = 0, 1, \dots, n)$ are in \mathbf{Z}_p . Then $f(p) = p/x + R$ and $f(x) = 1/p + R$. Suppose that K/R is injective. Then there exists an R -homomorphism $g: R \rightarrow K/R$ such that $g(a) = f(a)$ for all $a \in I$. We put $g(1) = k + R (k \in K)$. Then $g(p) = pk + R = p/x + R = f(p)$ and $g(x) = xk + R = 1/p + R = f(x)$. Since $pk - p/x \in R$ and $xk - x/p \in R$, $k =$

$$\frac{p + c_0x + c_1x^2 + \dots + c_mx^{m+1}}{px} = \frac{1 + pd_0 + pd_1x + \dots + pd_nx^n}{px}$$

for some $c_i \in \mathbf{Z}_p (i = 0, 1, \dots, m)$ and $d_j \in \mathbf{Z}_p (j = 0, 1, \dots, n)$. Thus $1 + pd_0 = p$ and so $d_0 = (p-1)/p$ does not belong to \mathbf{Z}_p . This is a contradiction. Hence K/R is not injective.

If a module M is nonsingular, then M^A has no nonzero injective submodule if and only if M has no nonzero injective submodule, where M^A is a direct product of copies of M for an index set A [2, Theorem 2.9]. But this is not true for some singular module M .

Lemma 7. *Let R be a commutative integral domain with quotient field $K \neq R$. Then the following assertions hold.*


- (1) K is an injective R -module.
- (2) $(K/R)^{R-(0)}$ has a nonzero injective submodule.

Proof. (1). Since K is divisible and is in $F(\sigma)$, it is injective. (2). We consider a correspondence $\phi: K \rightarrow (K/R)^{R-(0)}$ defined by $\phi(k) = (\dots, k/r_\alpha, \dots)$. Then ϕ is an R -homomorphism and $\text{Ker}(\phi) = \{k \in K \mid k/r_\alpha \in R \text{ for all } r_\alpha \in R - \{0\}\}$. Clearly ϕ is a monomorphism. By (1), $(K/R)^{R-(0)}$ has a nonzero injective submodule.

Example 8. *Let $R = \mathbf{Z}_p + x\mathbf{Q}[[x]]$, where $\mathbf{Q}[[x]]$ is the ring of formal power series over \mathbf{Q} and K the quotient field of R . Then K/R has no nonzero*

injective submodule but $(K/R)^{R^{-(0)}}$ has a nonzero injective submodule.

Proof. It is sufficient to show that K/R is indecomposable and it is not injective. First we show that K/R is indecomposable. We assume that $K/R = A/R \oplus B/R$, where A and B are R -submodules of K containing R with $K = A+B$ and $A \cap B = R$. We claim that $1/x$ belongs either to A or to B . Since $1/x$ is in $A+B$, there exist $\alpha \in A$ and $\beta \in B$ such that $1 = \alpha x + \beta x$. Thus $\alpha x = 1 - \beta x$ is in B and so αx is in R . Similarly βx is in R . Therefore $\alpha x = a_0 + a_1x + \cdots + a_nx^n + \cdots$ and $\beta x = b_0 + b_1x + \cdots + b_nx^n + \cdots$ for some a_0 and b_0 is \mathbf{Z}_p and a_i and b_i ($i = 1, 2, \cdots$) in \mathbf{Q} . Since $1 = \alpha x + \beta x$, $a_0 + b_0 = 1$ and $a_i + b_i = 0$ for $i \geq 1$. Thus either a_0 or b_0 is a unit. If a_0 (resp. b_0) is a unit, then αx (resp. βx) is unit in R and so $1/x = \alpha(\alpha x)^{-1}$ (resp. $\beta(\beta x)^{-1}$) is in A (resp. B). Thus we may assume that $1/x$ is in A . Then $\mathbf{Q}[[x]]$ is an R -submodule of A . In fact take $\gamma = c_0 + c_1x + c_2x^2 + \cdots$ be in $\mathbf{Q}[[x]]$. If $c_0 = 0$, then γ is in $R \subset A$. On the other hand, if $c_0 \neq 0$, then $c_0 = (1/x) \cdot c_0x \in A$. Since $c_1x + c_2x^2 + \cdots$ is in R , γ is in A and so $\mathbf{Q}[[x]]$ is an R -submodule of A . Next we show that a/x belongs to A for every $a \in A$. Indeed, let $a/x = a' + \beta'$ ($a' \in A$ and $\beta' \in B$). Then $a = xa' + x\beta'$ and so $x\beta' = a - xa'$. Thus $x\beta'$ is in $A \cap B = R$. We put $x\beta' = d_0 + d_1x + d_2x^2 + \cdots$, where $d_0 \in \mathbf{Z}_p$ and $d_i \in \mathbf{Z}$ ($i = 1, 2, \cdots$). Then $\beta' = c_0/x + \sum_{i=1}^{\infty} d_ix^{i-1}$. Since c_0/x is in A and $\sum_{i=1}^{\infty} d_ix^{i-1}$ is in $\mathbf{Q}[[x]]$, β' is in A . Hence a/x is in A . As is easily seen, each element of K is of the form $h(x)/x^m$, where $h(x)$ is in $\mathbf{Q}[[x]]$ and m is a non-negative integer. Thus $K = A$, namely K/R is indecomposable. Secondly we show that K/R is not injective. We put $I = x\mathbf{Q}[[x]]$ and $I_n = (x/p^n)R$ for $n = 0, 1, 2, \cdots$. Then we have an infinite ascending chain $I_0 \subsetneq I_1 \subsetneq I_2 \subsetneq \cdots \subsetneq I_n \subsetneq I_{n+1} \subsetneq \cdots$ with $\bigcup_{n=0}^{\infty} I_n = I$. Note that any homomorphism φ_n from I_n to K/R is uniquely determined by an element α_n of K/R which can be arbitrarily chosen and by the equation $\varphi_n(a) = a\alpha_n$ for all $a \in I_n$. Let $\{a_i\}$ be a sequence of integers such that $0 \leq a_i < p$ for all $i \geq 0$. For each n , let $f_n = (1/x)\sum_{i=0}^n a_i p^i$ be an element of K and $\alpha_n = f_n + R$ an element of K/R . Then we have $\alpha_{n+1} - \alpha_n = (1/x)a_{n+1}p^{n+1} + R$. Since $a_{n+1}p$ is in \mathbf{Z} , $(x/p^n)(\alpha_{n+1} - \alpha_n) = 0$ for $n = 0, 1, 2, \cdots$. Let φ_n be the R -homomorphism from I_n to K/R corresponding to α_n as noted above. Then the above equations imply that the system (I_n, φ_n) forms a direct system and $\varinjlim I_n = I$, we have the following commutative diagram



where $\varphi(a) = \varphi_n(a) = a\alpha_n$ for all $a \in I_n$.

Assume that K/R is injective. Then there exists an element α of K/R such that $\varphi(s) = s\alpha$ for all $s \in I$. If we take $s = x/p^n \in I_n$, then $\varphi(s) = s\alpha = s\alpha_n = \varphi_n(s)$ and so $(x/p^n)(\alpha - \alpha_n) = 0$. If we put $\alpha = f + R$, where $f \in K$, it follows that $(x/p^n)(f - f_n) \in R$ for all n , namely, $xf - \sum_{i=0}^n a_i p^i \in p^n R$. Hence we have xf is in R . We put $xf = \sum_{i=0}^{\infty} b_i p^i$, where $b_0 \in \mathbb{Z}_p$ and $b_i \in \mathbb{Q}$ for $i \geq 1$. Then we have $b_0 = \sum_{i=0}^n a_i p^i + p^n c_0$ for some $c_0 \in \mathbb{Z}_p$. Thus the sequence $\{\sum_{i=0}^n a_i p^i\}_n$ converges to b_0 with respect to the $p\mathbb{Z}_p$ -adic topology on \mathbb{Z}_p , that is, the p -adic number $\sum_{i=0}^{\infty} a_i p^i$ represents a rational number. This is absurd because the sequence $\{a_i\}_i$ of integers can be arbitrarily chosen. It follows that K/R is not injective.

Acknowledgements. The authors wish to thank to Professor A. Facchini for his helpful suggestions about example 8.

REFERENCES

- [1] S. MORIMOTO: On left exact preradicals, *Math. J. Okayama Univ.* **27** (1985), 87—95.
- [2] S. MORIMOTO: On some type of left exact radicals, *Math. J. Okayama Univ.* **32** (1990), 129—138.
- [3] B. STENSTROM: *Rings of Quotients*, Grundle. Math. Wiss. 217, Springer Verlag Berlin, 1975.

S. MORIMOTO
 HAGI KOEN GAKUIN HIGH SCHOOL
 HIGASHITAMACHI, HAGI, YAMAGUCHI, JAPAN 758

T. SHUDO
 DEPARTMENT OF GENERAL EDUCATION
 FUKUOKA DENTAL COLLEGE
 SAWARA-KU, FUKUOKA 814-01, JAPAN

(Received October 18, 1991)