

## **$K$ -SEMIMETRIZABILITIES AND $C$ -STRATIFIABILITIES OF SPACES**

IWAO YOSHIOKA

### 1. INTRODUCTION AND DEFINITIONS

In 1966, Arhangel'skiĭ [1] introduced the concepts of symmetrizable spaces and he showed that a  $T_2$ -space is metrizable if, and only if, it has a compatible symmetric  $d$  satisfying condition (A):  $d(F, K) > 0$  for any disjoint closed subset  $F$  and compact subset  $K$ . Also, Arhangel'skiĭ gave the class of spaces with a compatible symmetric  $d$  satisfying condition (K):  $d(H, K) > 0$  for any disjoint compact subsets  $H$  and  $K$ , and he conjectured that every symmetrizable space has a compatible symmetric satisfying condition (K). After that, in 1975, Martin [26] presented the question on whether every regular semimetrizable space is  $K$ -semimetrizable (i.e. it has a compatible symmetric satisfying condition (K)), or if every Moore space is  $K$ -semimetrizable. In 1979, Burke [6] gave a negative answer that there exists a separable Moore space which is not  $K$ -semimetrizable.

Lee [22] defined the class of  $c$ -stratifiable spaces which contains the classes of spaces with a regular  $G_\delta$ -diagonal and of  $\gamma$ ,  $T_2$ -spaces. He proved that a space  $X$  is  $K$ -semimetrizable if, and only if,  $X$  is  $c$ -stratifiable semimetrizable if, and only if,  $X$  is regular  $c$ -stratifiable, first countable and  $\beta$ . On the other hand, in [31], we introduced the concepts of strong  $\alpha$ -ness and showed that every strongly  $\alpha$ ,  $wM$ -space is metrizable. The properties of strongly  $\alpha$ -spaces were also studied in the same paper.

In this note, we study the relations among  $c$ -stratifiable spaces, strongly  $\alpha$ -spaces,  $K$ -semimetrizable spaces, developable spaces and Nagata spaces, and the conditions for spaces to be  $K$ -semimetrizable or full  $K$ -semimetrizable.

We prove that a space  $X$  is  $K$ -semimetrizable if, and only if, it is a  $c$ -stratifiable  $q$ ,  $\beta$ -space. We also show that a space  $X$  is full  $K$ -semimetrizable if, and only if, it is a  $w\theta$ ,  $\beta$ -space with a regular  $G_\delta$ -diagonal, which is a slight generalization of [32; Theorem 2]. We also show that a space  $X$  is Nagata if, and only if, it is  $K$ -semimetrizable  $wcc$  if, and only if, it is regular semimetrizable  $wcc$ . Moreover, for metrizations of  $wM$ -spaces, we have that every  $wM$ -space with a  $G_\delta^*$ -diagonal is metrizable.

In §2, we study the relations between  $c$ -stratifiable spaces and strongly  $\alpha$ -spaces. Also, we consider the conditions for spaces to be strongly  $\alpha$  or  $c$ -stratifiable. In particular, we show that in the realm of  $c$ -stratifiable spaces,  $wN$ -spaces are Nagata,  $q$ -spaces are first countable,  $wcc$ -spaces are  $k$ -semistratifiable and  $w\Delta$ -spaces are developable.

In §3, we study the class of  $K$ -semimetrizable spaces. First, we show that a space  $X$  is  $K$ -semimetrizable if, and only if, it is  $c$ -stratifiable  $q, \beta$ . Secondly, we prove that in the class of pseudocompact spaces or locally connected rim-compact spaces, developable  $K$ -semimetrizable spaces are equivalent to  $c$ -stratifiable  $\beta$ -spaces (or  $K$ -semimetrizable spaces), and every metacompact  $p$ -space with a  $G_\delta$ -diagonal is a  $K$ -semimetrizable Moore space.

In §4, for the class of  $w\theta, wcc$ -spaces which contains the class of  $wM$ -spaces, we show that every  $w\theta, wcc$ -space with a  $G_\delta^*$ -diagonal is metrizable and every  $c$ -stratifiable  $w\theta, wcc$ -space is metrizable.

Throughout this paper, we assume that all spaces are  $T_1$ , but paracompactness is assumed to be  $T_2$ . We denote a sequence  $\{x_n | n \in \mathbb{N}\}$  by  $\{x_n\}$  and the set of natural numbers by  $\mathbb{N}$ . Finally, we refer the reader to [9] for undefined terms.

**Definition 1.1.** A  $g$ -function on a space  $X$  with a topology  $\mathcal{T}$  is a map  $g : \mathbb{N} \times X \rightarrow \mathcal{T}$  such that  $g(n, x) = g_n(x)$  is an open neighbourhood of  $x$  for every  $x \in X$  and each  $n \in \mathbb{N}$  and we denote the map  $g$  by  $(\{g_n(x) | x \in X\})$ . For a subset  $A$  of  $X$ , we put  $g_n(A) = \cup\{g_n(x) | x \in A\}$ .

A point  $p$  in  $X$  is called a *cluster point* of a sequence  $\{x_n\} \subset X$  if any open neighbourhood of  $p$  contains  $x_n$  for infinitely many  $n$ 's.

For a space  $X$ , we now consider the following conditions on a  $g$ -function  $(\{g_n(x) | x \in X\})$ .

- (A) If  $g_n(x) \cap g_n(x_n) \neq \emptyset$  ( $n \geq 1$ ), then  $x$  is a cluster point of  $\{x_n\}$ .
- (B) If  $g_n(x) \cap g_n(x_n) \neq \emptyset$  ( $n \geq 1$ ), then  $\{x_n\}$  has a cluster point.
- (C) If  $x \in g_n(x_n)$  ( $n \geq 1$ ), then  $x$  is a cluster point of  $\{x_n\}$ .
- (D) If  $x \in g_n(x_n)$  ( $n \geq 1$ ), then  $\{x_n\}$  has a cluster point.
- (E) If  $y_n \in g_n(x_n)$  ( $n \geq 1$ ) and  $\{y_n\}$  has a cluster point, then  $\{x_n\}$  has a cluster point.
- (F) If  $x_n \in g_n(x)$  ( $n \geq 1$ ), then  $\{x_n\}$  has a cluster point.
- (G) If  $y_n \in g_n(p), x_n \in g_n(y_n)$  ( $n \geq 1$ ), then  $p$  is a cluster point of  $\{x_n\}$ .
- (H) If  $y_n \in g_n(p), x_n \in g_n(y_n)$  ( $n \geq 1$ ), then  $\{x_n\}$  has a cluster point.
- (I) If  $y_n \in g_n(p), x_n, p \in g_n(y_n)$  ( $n \geq 1$ ), then  $p$  is a cluster point of  $\{x_n\}$ .
- (J) If  $y_n \in g_n(p), x_n, p \in g_n(y_n)$  ( $n \geq 1$ ), then  $\{x_n\}$  has a cluster point.
- (K) If  $x_n, p \in g_n(y_n)$  ( $n \geq 1$ ), then  $p$  is a cluster point of  $\{x_n\}$ .
- (L) If  $x_n, p \in g_n(y_n)$  ( $n \geq 1$ ), then  $\{x_n\}$  has a cluster point.

In the above conditions (A)-(L), we can assume that  $g_{n+1}(x) \subset g_n(x)$  for every  $x \in X$  and each  $n \in \mathbb{N}$ .

**Definition 1.2.** A space with a  $g$ -function satisfying (A) is called a *Nagata space* [15] (Nagata spaces were first defined by Ceder [7]) and a space with a  $g$ -function satisfying (B) is called a  $wN$ -space [18]. In this case the  $g$ -function is called a *Nagata-function* (a  $wN$ -function, respectively).

**Definition 1.3.** A space  $X$  is called a *semistratifiable* ( $\beta$ -,  $wcc$  (=weak contraconvergent)-,  $q$ -,  $\gamma$ -,  $w\gamma$ -,  $\theta$ -,  $w\theta$ -) space if  $X$  has a  $g$ -function satisfying (C) ( (D), (E), (F), (G), (H), (I), (J), respectively). (See [17], [18] and [31])

The following result is not difficult to see.

**Proposition 1.4.** [31; Theorem 3.5] *A space  $X$  is  $wN$  if, and only if, it is  $q$  and  $wcc$ .*

**Definition 1.5.** A space  $X$  is called *stratifiable* [3] (equivalently,  $M_3$  [7]) if  $X$  has a  $g$ -function that satisfies (C) and if  $x \notin \overline{g_m(F)}$  for some  $m \in \mathbb{N}$ , whenever  $F$  is closed and  $x \notin F$ . The class of  $k$ -semistratifiable spaces introduced by Lutzer [24] can be characterized by the following conditions [12, 31]. A space  $X$  is  $k$ -semistratifiable if, and only if,  $X$  has a  $g$ -function ( $\{g_n(x)\} | x \in X$ ) such that  $g_m(F) \cap K = \emptyset$  for some  $m \in \mathbb{N}$ , whenever  $F$  is closed,  $K$  is compact and  $F \cap K = \emptyset$ , if, and only if, in the class of  $T_2$ -spaces,  $X$  has a  $g$ -function ( $\{g_n(x)\} | x \in X$ ) such that whenever  $y_n \in g_n(x_n)$  ( $n \geq 1$ ) and  $\{y_n\} \rightarrow y$ , then  $\{x_n\} \rightarrow y$ .

The following implications are known.

Nagata  $\implies$  stratifiable  $\implies k$ -semistratifiable  $\implies$  semistratifiable  $\implies \beta$ .

Also, it is known that a Nagata space is equivalent to a first countable stratifiable space and every stratifiable space is paracompact. Every semistratifiable space  $X$  is subparacompact and has a  $G_\delta$ -diagonal if it is  $T_2$  [14; Theorem 5.11].

## 2. C-STRATIFIABLE SPACES AND STRONGLY $\alpha$ -SPACES

We begin by considering the relations between  $c$ -stratifiable spaces and strongly  $\alpha$ -spaces, and the conditions for spaces to be  $c$ -stratifiable or strongly  $\alpha$ .

**Definition 2.1.** A space  $X$  is called  *$c$ -stratifiable* [22] ( *$c$ -semistratifiable* [25]) if  $X$  has a  $g$ -function such that if  $x \notin K$ , where  $K$  is compact, then  $x \notin \overline{g_m(K)}$  ( $x \notin g_m(K)$ ; in [25], it is assumed that  $K$  is closed compact) for some  $m \in \mathbb{N}$ . A space  $X$  is called  *$cs$ -stratifiable* if  $X$  has a  $g$ -function such

that if  $x \notin C$ , where  $C$  is the union of a convergent sequence and any one of its limit points, then  $x \notin \overline{g_m(C)}$  for some  $m \in \mathbb{N}$ . A space  $X$  is called *weak  $c$ -stratifiable* if  $X$  has a  $g$ -function such that, whenever  $C$  and  $D$  are disjoint compact subsets, then  $g_m(C) \cap D = \emptyset$  for some  $m \in \mathbb{N}$ .

Every stratifiable space or  $\gamma$ ,  $T_2$ -space is  $c$ -stratifiable [22], but the Sorgenfrey line is a paracompact  $c$ -stratifiable space which is not semistratifiable. Also, every  $c$ -stratifiable space is  $cs$ -stratifiable and weak  $c$ -stratifiable [22; Theorem 1.3], every  $cs$ -stratifiable space is  $T_2$  and every semistratifiable  $T_2$ -space is  $c$ -semistratifiable.

**Definition 2.2.** A space  $X$  is called *strongly  $\alpha$*  [31] ( $\alpha$  [17]) if  $X$  has a  $g$ -function such that (i) for each  $n \in \mathbb{N}$ ,  $y \in g_n(x) \implies g_n(y) \subset g_n(x)$  and (ii)  $\bigcap_{n \geq 1} \overline{g_n(x)} = \{x\}$  ( $\bigcap_{n \geq 1} g_n(x) = \{x\}$ ).

For  $g$ -functions in Definitions 2.1 and 2.2, we can assume that  $g_{n+1}(x) \subset g_n(x)$  for every  $x \in X$  and each  $n \in \mathbb{N}$ . Every strongly  $\alpha$ -space is also  $T_2$ .

**Theorem 2.3.** *Every  $cs$ -stratifiable  $q$ -space  $X$  or regular weak  $c$ -stratifiable  $q$ -space  $X$  is a first countable  $c$ -stratifiable space.*

*Proof.* Let  $g$  be a  $q$  and  $cs$ -stratifiable function of a space  $X$ . We show that  $\{g_n(x)\}$  is an open neighbourhood base of  $x$  for every  $x \in X$ . Suppose that  $x \in X$  and  $x_n \in g_n(x) \setminus U$  ( $n \geq 1$ ) for some open neighbourhood  $U$  of  $x$ . Since  $g$  is a  $q$ -function,  $\{x_n\}$  has a cluster point  $p$  and  $p \notin \{x\}$ . Since  $g$  is a  $cs$ -stratifiable function,  $p \notin \overline{g_m(x)} \supset \overline{\{x_j | j \geq m\}} \ni p$  for some  $m \in \mathbb{N}$ . This contradiction implies that  $\{g_n(x)\}$  is a neighbourhood base of  $x$ . To see that  $g$  is a  $c$ -stratifiable function, suppose that  $x \notin K$ , where  $K$  is compact in  $X$ , and  $x \in \bigcap_{n \geq 1} \overline{g_n(K)}$ . Then, there exist sequences  $\{x_n\} \subset K$  and  $\{y_n\}$  such that  $y_n \in g_n(x) \cap g_n(x_n)$ . Since  $K$  is sequentially compact,  $\{x_{n(i)}\} \longrightarrow p$  for some point  $p \in K$  and some subsequence  $\{x_{n(i)}\} \subset \{x_n\}$ , and  $\{y_{n(i)}\} \longrightarrow x$  for the subsequence  $\{y_{n(i)}\} \subset \{y_n\}$ . Then  $x \notin \{x_{n(i)} | i \geq 1\} \cup \{p\} = C$ , and hence  $x \notin \overline{g_m(C)}$  for some  $m \in \mathbb{N}$ . Therefore, for some  $n(j) \geq m$ ,  $y_{n(j)} \notin \overline{g_m(C)} \supset \overline{g_{n(j)}(x_{n(j)})} \ni y_{n(j)}$ . This contradiction implies that  $g$  is also a  $c$ -stratifiable function.

For the second part, we can assume that  $g$  is a weak  $c$ -stratifiable  $q$ -function satisfying  $g_{n+1}(x) \subset g_n(x)$ . If  $x$  and  $y$  are distinct points, then  $g_m(x) \cap \{y\} = \emptyset$  for some  $m \in \mathbb{N}$ . Hence,  $\bigcap_{n \geq 1} \overline{g_n(x)} = \{x\}$ . Suppose that  $x \in X$  and  $x_n \in g_n(x) \setminus U$  ( $n \geq 1$ ) for some open neighbourhood  $U$  of  $x$ . Then  $\{x_n\}$  has a cluster point  $p$ . Hence,  $p \in \overline{g_n(x)}$  for each  $n \in \mathbb{N}$ . This contradiction asserts that  $\{g_n(x)\}$  is a neighbourhood base of  $x$ . Therefore,

$g$  is a  $c$ -stratifiable function by [22; Theorem 1.3].

**Theorem 2.4.** (1) *Every strongly  $\alpha$ -space  $X$  or  $k$ -semistratifiable  $T_2$ -space  $X$  is weak  $c$ -stratifiable.*

(2) *Every strongly  $\alpha$ ,  $q$ -space  $X$  is  $c$ -stratifiable.*

*Proof.* (1): First, let  $g$  be a strongly  $\alpha$ -function of  $X$ . Suppose that there are disjoint compact subsets  $C$  and  $D$  such that  $x_n \in g_n(C) \cap D$  ( $n \geq 1$ ). Then  $x_n \in g_n(y_n)$  for some sequence  $\{y_n\} \subset C$ . Hence  $\{y_n\}$  clusters at a point  $y \in C$  and contains a subsequence  $\{y_{n(i)}\}$  such that  $y_{n(i)} \in g_i(y)$ . Then  $x_{n(i)} \in g_{n(i)}(y_{n(i)}) \subset g_i(y_{n(i)}) \subset g_i(y)$  ( $i \geq 1$ ), and  $\{x_{n(i)}\}$  has a cluster point  $x \in D$ . Then for each  $i \in \mathbb{N}$ ,  $x \in \overline{\{x_{n(j)} \mid j \geq i\}} \subset \overline{g_i(y)}$ . Therefore  $x = y$ , which is a contradiction. Next, that a  $k$ -semistratifiable  $T_2$ -space is weak  $c$ -stratifiable follows from the equivalent condition of a  $k$ -semistratifiable space in Definition 1.5.

(2): Let  $g$  be a  $q$ -function and  $h$  be a strongly  $\alpha$ -function of a space  $X$ . Here, we can assume that  $g_n(x) \subset h_n(x)$ . For some  $x \in X$  and some compact subset  $K$ , suppose that  $x \notin K$  and  $x \in \bigcap_{n \geq 1} \overline{g_n(K)}$ . Then there exist sequences  $\{y_n\}$  and  $\{z_n\}$  such that  $y_n \in K$  and  $z_n \in g_n(x) \cap g_n(y_n)$ . Let  $y \in K$  be a cluster point of  $\{y_n\}$ . Then  $y_{n(i)} \in g_i(y)$  for some increasing subsequence  $\{n(i)\}$  of  $\mathbb{N}$ . Also, since  $z_{n(i)} \in g_i(x)$  ( $i \geq 1$ ),  $\{z_{n(i)}\}$  has a cluster point  $z$ . Since  $y_{n(i)} \in h_i(y)$  ( $i \geq 1$ ),  $z_{n(i)} \in g_{n(i)}(y_{n(i)}) \subset h_i(y_{n(i)}) \subset h_i(y)$ . Therefore,  $\{z_{n(j)} \mid j \geq i\} \subset h_i(y)$  ( $i \geq 1$ ) and hence,  $z \in \overline{h_i(y)}$  ( $i \geq 1$ ), which implies that  $y = z$ . Moreover, since  $z_{n(i)} \in g_i(x) \subset h_i(x)$  ( $i \geq 1$ ), we have  $\{z_{n(j)} \mid j \geq i\} \subset h_i(x)$ , and hence  $z \in \overline{h_i(x)}$ . Consequently,  $x = z$ . This contradiction implies that  $g$  is a  $c$ -stratifiable function.

We now study the conditions for spaces to be  $c$ -stratifiable or strongly  $\alpha$ .

**Definition 2.5.** A space  $X$  is called a  $w\Delta$ -space [4] if it has a sequence  $\{\mathcal{G}_n\}$  of open covers such that whenever  $x_n \in st(x, \mathcal{G}_n)$  ( $n \geq 1$ ), then  $\{x_n\}$  has a cluster point. A space  $X$  is called a *developable* space if it has a sequence  $\{\mathcal{G}_n\}$  of open covers such that for each  $x \in X$ , the sequence  $\{st(x, \mathcal{G}_n)\}$  is a neighbourhood base of  $x$ . A regular developable space is called a *Moore* space. These spaces are characterized by  $g$ -functions as follows [18]: A space  $X$  is  $w\Delta$  (*developable*) if and only if  $X$  has a  $g$ -function satisfying (L) ((K), respectively).

**Definition 2.6.** (1) For each  $k \in \mathbb{N}$ , a space  $X$  is said to have a  $G_\delta(k)$ -diagonal if  $X$  has a sequence  $\{\mathcal{G}_n\}$  of open covers such that for any distinct points  $x$  and  $y$ , there exists  $m \in \mathbb{N}$  such that  $y \notin st^k(x, \mathcal{G}_m)$ , where  $st^{k+1}(x, \mathcal{G}_m) = st(st^k(x, \mathcal{G}_m))$ .

(2) A sequence  $\{\mathcal{G}_n\}$  of open covers of a space  $X$  is said to satisfy the 3-link property [32] (equivalently, it is a  $G_\delta(3)$ -diagonal sequence) if it is true that for any distinct points  $x$  and  $y$ , there exists  $m \in \mathbb{N}$  such that no member of  $\mathcal{G}_m$  intersects both  $st(x, \mathcal{G}_m)$  and  $st(y, \mathcal{G}_m)$ .

(3) A space  $X$  is said to have a regular  $G_\delta$ -diagonal [32] if there is a sequence  $\{\mathcal{G}_n\}$  of open covers of  $X$  such that if  $x$  and  $y$  are distinct points of  $X$ , then there are an integer  $m$  and open neighbourhoods  $U$  and  $V$  of  $x$  and  $y$ , respectively, such that no member of  $\mathcal{G}_m$  intersects both  $U$  and  $V$ .

(4) A space  $X$  is said to have a  $G_\delta^*$ -diagonal if  $X$  has a sequence  $\{\mathcal{G}_n\}$  of open covers such that whenever  $x \neq y$ , there exists  $m \in \mathbb{N}$  that satisfies  $y \notin st(x, \mathcal{G}_m)$ .

It is easily seen that for a sequence  $\mathcal{G} = \{\mathcal{G}_n\}$  of open covers of a space  $X$ ,  $\mathcal{G}$  is  $G_\delta(2)$ -diagonal if, and only if, whenever  $x \neq y$ , there exists  $m \in \mathbb{N}$  satisfying  $x \notin st(p, \mathcal{G}_m)$  or  $y \notin st(p, \mathcal{G}_m)$  for every  $p \in X$  (this property is called *strong  $G_\delta$ -diagonal* in [31]).

We note that for properties of a sequence  $\{\mathcal{G}_n\}$  of open covers of a space  $X$ , the following implications hold:

3-link property  $\Rightarrow$  regular  $G_\delta$ -diagonal  $\Rightarrow G_\delta^*$ -diagonal  $\Rightarrow G_\delta$ -diagonal and  
 3-link property  $\Rightarrow G_\delta(2)$ -diagonal = strong  $G_\delta$ -diagonal  $\Rightarrow G_\delta^*$ -diagonal.

In the realm of paracompact spaces, these properties are all equivalent. Every Nagata space is paracompact and has a  $G_\delta$ -diagonal. Every developable  $T_2$ -space has a  $G_\delta(2)$ -diagonal and every regular semistraifiable space has a  $G_\delta^*$ -diagonal [14, 17]. On the other hand, the space  $\Psi$  in Example 4.5 is a Moore space which does not have a regular  $G_\delta$ -diagonal.

**Definition 2.7.** (1) A space  $X$  is called *orthocompact* if every open cover of  $X$  has an open refinement  $\mathcal{V}$  such that  $\cap \mathcal{W} = \cap \{W \mid W \in \mathcal{W}\}$  is open for every  $\mathcal{W} \subset \mathcal{V}$ .

(2) A space  $X$  is called *submetrizable* if there is a continuous one-to-one map from  $X$  onto a metric space.

It is well known that the following implications hold:  
 metacompact spaces  $\implies$  orthocompact spaces, and  
 stratifiable spaces  $\implies$  paracompact spaces with a  $G_\delta$ -diagonal [3, 29]  $\implies$   
 submetrizable spaces .

**Theorem 2.8.** (1) *Every space  $X$  with a regular  $G_\delta$ -diagonal is  $c$ -stratifiable.*

- (2) Every orthocompact space  $X$  with a  $G_\delta^*$ -diagonal, or orthocompact regular space  $X$  with a  $G_\delta$ -diagonal is strongly  $\alpha$ .
- (3) Every orthocompact developable  $T_2$ -space  $X$  is strongly  $\alpha$  and  $c$ -stratifiable.
- (4) Every orthocompact regular  $c$ -semistratifiable  $\beta$ -space  $X$  is strongly  $\alpha$ .
- (5) Every submetrizable space  $X$  is strongly  $\alpha$  and  $c$ -stratifiable.

*Proof.* (1) is proved in [22; Proposition 3.2].

(2): Let  $X$  be a regular space and let  $\{\mathcal{G}_n\}$  be a  $G_\delta$ -diagonal sequence of  $X$ . Then for each  $n \in \mathbb{N}$ ,  $\mathcal{G}_n$  has an open refinement  $\mathcal{H}_n$  such that  $\{\overline{H} | H \in \mathcal{H}_n\}$  is a refinement of  $\mathcal{G}_n$  and  $\cap \mathcal{W}$  is open for every  $\mathcal{W} \subset \mathcal{H}_n$ . Therefore, in both cases, we may assume that there exists a sequence  $\{\mathcal{H}_n\}$  of open covers such that

- (i) for each  $n \in \mathbb{N}$ ,  $\cap \mathcal{W}$  is open for every  $\mathcal{W} \subset \mathcal{H}_n$ , and
- (ii) for distinct points  $x$  and  $y$ , there is an  $m \in \mathbb{N}$  such that  $x \in H \subset \overline{H}$  and  $y \notin \overline{H}$  for some  $H \in \mathcal{H}_m$ .

Here, for any  $x \in X$  and each  $n \in \mathbb{N}$ , we put  $h_n(x) = \cap \{H \in \mathcal{H}_n | x \in H\}$ . Then the  $g$ -function  $(\{h_n(x) | x \in X\})$  satisfies the conditions of Definition 2.2.

(3): Since every developable  $T_2$ -space has a  $G_\delta^*$ -diagonal,  $X$  is a strongly  $\alpha$  from (2) and it is  $c$ -stratifiable from Theorem 2.4.

(4): Since every regular  $c$ -semistratifiable  $\beta$ -space is semistratifiable [25; Theorem 3], it has a  $G_\delta$ -diagonal. Hence  $X$  is strongly  $\alpha$  from (2).

(5): Let  $f : X \rightarrow M$  be a continuous one-to-one onto map, where  $M$  is a metric space. By (3),  $M$  is strongly  $\alpha$  and  $c$ -stratifiable. Therefore (5) follows from the following fact:

*Let  $f : X \rightarrow Y$  be a continuous one-to-one onto map. If  $h$  is a strongly  $\alpha$ -function ( $c$ -stratifiable function) of  $Y$ , then  $(\{g_n(x) | x \in X\})$ , where  $g_n(x) = f^{-1}[h_n(f(x))]$ , is a strongly  $\alpha$ -function ( $c$ -stratifiable function, respectively) of  $X$ .*

We note that every metacompact regular semistratifiable  $q$ -space is strongly  $\alpha$  and hence it is  $c$ -stratifiable. Also, every regular  $k$ -semistratifiable  $q$ -space is Nagata [31], and hence it is strongly  $\alpha$ ,  $c$ -stratifiable. On the other hand, the separable Moore space  $X$  in Example 4.6 is neither strongly  $\alpha$  nor  $c$ -stratifiable.

The following question arises naturally from (4) of the above Theorem.

**Question 2.9.** Is every paracompact (or metacompact regular)  $c$ -semistratifiable  $q$ -space,  $c$ -stratifiable ?

**Theorem 2.10.** *For a cs-stratifiable space  $X$ , the following implications hold:*

(1)  $\beta \implies$  *semistratifiable*, (2)  $wcc \implies$   *$k$ -semistratifiable*, (3)  $wN \implies$  *Nagata*, (4)  $w\theta \implies \theta$ , (5)  $w\gamma \implies \gamma$  and (6)  $w\Delta \implies$  *developable*.

*Proof.* (1): Let  $g$  be a cs-stratifiable and  $\beta$ -function of  $X$  and let  $x \in g_n(x_n)$  ( $n \geq 1$ ). For any subsequence  $\{x_{n(i)}\}$  of  $\{x_n\}$ ,  $\{x_{n(i)}\}$  has a cluster point since  $x \in g_i(x_{n(i)})$  ( $i \geq 1$ ). Let  $p$  be a cluster point of  $\{x_n\}$  and  $p \neq x$ . Then  $\{x_n\}$  contains a subsequence  $S = \{x_{n(j)}\}$  such that  $x_{n(j)} \in g_j(p) \setminus \{x\}$  ( $j \geq 1$ ). Since  $\{x_{n(k)} \mid k \geq j\} \subset g_j(p)$  ( $j \geq 1$ ) and  $\{p\} = \bigcap_{j \geq 1} \overline{g_j(p)}$ ,  $p$  is the only cluster point of  $S$ . Hence  $S$  converges to  $p$ . Since  $x \notin C = S \cup \{p\}$ ,  $x \notin \overline{g_m(C)}$  for some  $m \in \mathbb{N}$ . But, for some  $n(k) \geq m$ ,  $x \in g_{n(k)}(x_{n(k)}) \subset g_m(C)$ . This contradiction implies that  $x = p$  is a cluster point of  $\{x_n\}$ .

(2): Let  $g$  be a cs-stratifiable,  $wcc$ -function satisfying  $\{x\} = \bigcap_{n \geq 1} \overline{g_n(x)}$  for every  $x \in X$ . Since  $X$  is a  $T_2$ -space, it is enough to show that  $g$  satisfies the  $k$ -semistratifiable condition of Definition 1.5. Let  $y_n \in g_n(x_n)$  ( $n \geq 1$ ) and  $\{y_n\} \longrightarrow y$ . First, we show that  $\{x_n\}$  contains a subsequence which converges to  $y$ . Indeed, for any subsequence  $\{x_{n(i)}\}$  of  $\{x_n\}$ ,  $y_{n(i)} \in g_{n(i)}(x_{n(i)}) \subset g_i(x_{n(i)})$  ( $i \geq 1$ ) and  $\{y_{n(i)}\} \longrightarrow y$ , hence  $\{x_{n(i)}\}$  has a cluster point. Let  $p$  be a cluster point of  $\{x_n\}$ . It is easily seen that there exists a subsequence  $S = \{x_{n(i)}\}$  of  $\{x_n\}$  such that  $x_{n(i)} \in g_i(p)$ . Since  $\overline{\{x_{n(j)} \mid j \geq i\}} \subset \overline{g_i(p)}$  for each  $i \in \mathbb{N}$ ,  $p$  is a unique cluster point of  $S$ . Hence  $S$  converges to  $p$ . If  $p \neq y$ , then  $y \notin \overline{\{x_{n(i)} \mid i \geq m\} \cup \{p\}} = C$  for some  $m \in \mathbb{N}$ . Therefore  $y \notin \overline{g_{n(k)}(C)}$  for some  $k \geq m$ . This contradiction asserts that  $S$  converges to  $y$ . Next, if  $\{x_n\}$  does not converge to  $y$ , then we have an open neighbourhood  $W$  of  $y$  and a subsequence  $\{x_{n(i)}\}$  such that  $\{x_{n(i)}\} \cap W = \emptyset$ . Then since  $y_{n(i)} \in g_{n(i)}(x_{n(i)}) \subset g_i(x_{n(i)})$  ( $i \geq 1$ ) and  $\{y_{n(i)}\} \longrightarrow y$ ,  $\{x_{n(i)}\}$  contains a subsequence which converges to  $y$ . This contradiction implies that  $\{x_n\} \longrightarrow y$ .

(3): Since  $X$  is  $q$  and  $wcc$ , by Theorem 2.3 and (2) of this theorem, there is a  $g$ -function  $g$  such that, whenever  $y_n \in g_n(x_n)$  ( $n \geq 1$ ) and  $\{y_n\} \longrightarrow y$ , then  $\{x_n\} \longrightarrow y$ , and  $\{g_n(x)\}$  is a neighbourhood base of  $x$ . To see that  $g$  is a Nagata function, let  $y_n \in g_n(x) \cap g_n(x_n)$  ( $n \geq 1$ ). Then  $\{y_n\} \longrightarrow x$ . Hence  $\{x_n\} \longrightarrow x$ .

(4): Let  $g$  be a cs-stratifiable  $w\theta$ -function of  $X$ . Then  $g$  is a  $q$ -function. Indeed, let  $x_n \in g_n(x)$  ( $n \geq 1$ ), then  $\{x_n\}$  has a cluster point since  $x \in g_n(x)$ ,  $x_n, x \in g_n(x)$  ( $n \geq 1$ ). Therefore,  $\{g_n(x)\}$  is a neighbourhood base of  $x$  by Theorem 2.3. Now, suppose that  $y_n \in g_n(p)$  and  $x_n, p \in g_n(y_n)$  ( $n \geq 1$ ). Then  $\{x_n\}$  has a cluster point  $x$  and  $\{y_n\}$  converges to  $p$ . If  $x \neq p$ , then



$x \notin \{y_n | n \geq m\} \cup \{p\} = C$  for some  $m \in \mathbb{N}$ . Therefore  $x \notin \overline{g_k(C)}$  for some  $k \geq m$ . This contradiction implies  $x = p$ , and hence  $g$  is a  $\theta$ -function.

(5): Let  $g$  be a  $cs$ -stratifiable,  $w\gamma$ -function of a space  $X$ . Suppose that  $y_n \in g_n(p), x_n \in g_n(y_n)$  ( $n \geq 1$ ). Then  $\{x_n\}$  has a cluster point  $x$ , and  $\{y_n\}$  converges to  $p$  since  $g$  is a  $q$ -function. If  $p \neq x$ , then  $x \notin \{y_n | n \geq m\} \cup \{p\} = C$  for some  $m \in \mathbb{N}$ . Therefore  $x \notin \overline{g_k(C)}$  for some  $k \geq m$ , which is a contradiction.

(6): Let  $g$  be a  $cs$ -stratifiable  $w\Delta$ -function of  $X$ . Since  $g$  is a  $\beta$ -function,  $g$  is a semistratifiable function. Now, suppose that  $x_n, p \in g_n(y_n)$  ( $n \geq 1$ ). Then  $\{y_n\}$  converges to  $p$  and  $\{x_n\}$  has a cluster point  $x$ . If  $x \neq p$ , then  $x \notin \{y_n | n \geq m\} \cup \{p\} = C$  for some  $m \in \mathbb{N}$ . Hence  $x \notin \overline{g_k(C)}$  for some  $k \geq m$ . This contradiction implies that the function  $g$  satisfies condition (K).

**Remark 2.11.** (1) In the class of strongly  $\alpha$ -spaces, the implications (3)-(6) in Theorem 2.10 are true by Theorem 2.4, (1) follows from [17; Theorem 5.2] and (2) follows from [31; Proposition 4.7].

(2) In the class of weak  $c$ -stratifiable regular spaces, the implications (3)-(6) in Theorem 2.10 are true by Theorem 2.3. For the implications (1) and (2), let  $g$  be a  $g$ -function satisfying the respective conditions. Then, since  $\bigcap_{n \geq 1} \overline{g_n(x)} = \{x\}$ , (1) and (2) are also true by a similar argument to the proof of Theorem 2.10.

The following questions regarding  $c$ -stratifiable spaces and strongly  $\alpha$ -spaces are natural.

**Question 2.12.** When are  $c$ -stratifiability and strong  $\alpha$ -ness coincident ?

**Question 2.13.** Is every paracompact first countable  $c$ -stratifiable space strongly  $\alpha$  ?

### 3. K-SEMIMETRIZABLE SPACES

**Definition 3.1.** Let  $X$  be a space. Then a function  $d : X \times X \rightarrow \mathbb{R}$  is called a *semimetric* if (i)  $d(x, y) \geq 0$ , (ii)  $d(x, y) = 0 \iff x = y$  and (iii)  $d(x, y) = d(y, x)$ .  $X$  is called a *semimetrizable* space or  $X$  has a *compatible semimetric* if there exists a semimetric  $d$  on  $X$  such that for any subset  $M \subset X, x \in \overline{M} \iff d(x, M) = 0$ , or equivalently, for any  $x \in X$  and any open neighbourhood  $U$  of  $x, x \in \text{int}B(x; \epsilon) \subset B(x; \epsilon) \subset U$  for some  $\epsilon > 0$ ; where  $B(A; \delta) = \{y \in X | d(A, y) = \inf\{d(a, y) | a \in A\} < \delta\}$  for each  $\delta > 0$  and any subset  $A \subset X$  and  $B(x; \delta) = B(\{x\}; \delta)$ . Then, for a sequence  $\{x_n\}$  in a semimetrizable space  $(X, d), \lim_{n \rightarrow \infty} d(x, x_n) = 0$

$\iff \{x_n\} \longrightarrow x$  in  $X$ . A semimetrizable space  $X$  with a compatible semimetric  $d$  is *K-semimetrizable* [26] if  $d(H, K) = \inf\{d(x, y) | x \in H, y \in K\} > 0$  for any disjoint compact subsets  $H$  and  $K$ . In this situation,  $d$  is called a *K-semimetric* on  $X$ .

It is well known [8] that a space  $X$  is semimetrizable if, and only if, it is a first countable, semistratifiable space.

**Definition 3.2.** Let  $(X, d)$  be a semimetrizable space. For each  $n \in \mathbb{N}$ , we put  $\mathcal{G}_n = \{\text{int}B(x; \epsilon) | \delta B(x; \epsilon) < 1/n\}$ , where for subset  $A$  of  $X$ ,  $\delta A = \sup\{d(x, y) | x, y \in A\}$ . The semimetric  $d$  is said to be *full* if  $\mathcal{G}_n$  is a cover of  $X$  for each  $n \in \mathbb{N}$ , or equivalently, if  $d$  satisfies Arhangel'skii's condition (AN): At each point, there is a neighbourhood of arbitrarily small diameter [1]. A space  $X$  is called *full K-semimetrizable* if  $X$  has a compatible full  $K$ -semimetric.

Zenor investigated spaces with a regular  $G_\delta$ -diagonal and gave the following result.

**Theorem 3.3** ([32; Theorem 2]). *For a space  $X$ , the following conditions are equivalent.*

- (1)  $X$  has a development satisfying the 3-link property.
- (2)  $X$  is a  $w\Delta$ -space with a regular  $G_\delta$ -diagonal.
- (3)  $X$  has a compatible semimetric  $d$  satisfying
  - (I) If  $\{x_n\} \longrightarrow x$  and  $\{y_n\} \longrightarrow x$ , then  $\lim_{n \rightarrow \infty} d(x_n, y_n) = 0$ , and
  - (II) If  $\{x_n\} \longrightarrow x$ ,  $\{y_n\} \longrightarrow y$  and  $x \neq y$ , then there exist  $r > 0$  and  $m \in \mathbb{N}$  such that  $d(x_n, y_n) > r$  for each  $n \geq m$ .

In substance, the first part of the following theorem is proved in (1)  $\iff$  (3) of [32; Theorem 2] or [22; Lemma 5.3].

**Theorem 3.4.** (1) *For a space  $X$ , the following conditions are equivalent.*

- (i)  $X$  is a developable space.
  - (ii)  $X$  has a compatible full semimetric  $d$ .
  - (iii)  $X$  has a compatible semimetric  $d$  satisfying (I) of Theorem 3.3.
- (2) *A space  $X$  is developable  $T_2$  if, and only if, it is  $w\theta$ ,  $\beta$  and has a  $G_\delta^*$ -diagonal.*

*Proof of (2).* We only prove the “if” part. Every  $\beta$ -space with a  $G_\delta^*$ -diagonal is semistratifiable [17; Theorem 5.2] and every semistratifiable  $w\theta$ -space is  $w\Delta$  [18; Proposition 4.5]. Hence  $X$  is developable [17; Theorem 2.5].

For a semimetric space, we have the following characterization. A regular space  $X$  is semimetrizable if, and only if, it is a  $q, \beta$ -space with a  $G_\delta^*$ -diagonal.

Indeed, let  $g$  be a  $q$ -function and  $\{\mathcal{G}_n\}$  be a  $G_\delta^*$ -diagonal sequence of a space  $X$ . We put  $h_n(x) = g_n(x) \cap st(x, \mathcal{G}_n)$ , then  $\{h_n(x)\}$  is a neighbourhood base of  $x$ . Also,  $X$  is semistratifiable from the proof of Theorem 3.4(2). For the converse implication, see [17].

The following theorem improves the result [11; Proposition 2.7] or [22; Theorem 5.2] that a space  $X$  is  $K$ -semimetrizable if, and only if, it is  $c$ -stratifiable and semimetrizable.

**Theorem 3.5.** *For a space  $X$ , the following conditions are equivalent.*

- (1)  $X$  is a  $K$ -semimetrizable space.
- (2)  $X$  is a  $c$ -stratifiable semimetrizable space.
- (3)  $X$  is a  $cs$ -stratifiable  $q, \beta$ -space.
- (4)  $X$  has a compatible semimetric  $d$  satisfying (II) of Theorem 3.3.
- (5)  $X$  has a compatible semimetric  $d$  such that,  $x \notin \overline{B(K; 1/m)}$  for some  $m \in \mathbb{N}$ , whenever  $x \notin K$  and  $K$  is compact.

*Proof.* (1) $\implies$ (2) is proved in [22; Theorem 5.2] and (2) $\implies$ (3) is evident.

(3) $\implies$ (1): Let  $g$  be a  $cs$ -stratifiable  $q, \beta$ -function of  $X$ . Then by Theorems 2.3 and 2.10,  $g$  is a  $c$ -stratifiable and semistratifiable function, and  $\{g_n(x)\}$  is an open neighbourhood base of  $x$  for every  $x \in X$ . Now, we define  $d(x, x) = 0$  and  $d(x, y) = 1/\inf\{j | x \notin g_j(y) \text{ and } y \notin g_j(x)\}$  if  $x \neq y$ . By [22; Theorem 5.2],  $(X, d)$  is  $K$ -semimetrizable.

(1) $\implies$ (4): Let  $d$  be a compatible  $K$ -semimetric on  $X$ . Suppose that  $\{x_n\} \rightarrow x, \{y_n\} \rightarrow y$  and  $x \neq y$ . Since  $X$  is  $T_2$ , for some  $m \in \mathbb{N}$ ,  $H = \{x_n | n \geq m\} \cup \{x\}$  and  $K = \{y_n | n \geq m\} \cup \{y\}$  are disjoint compact subsets. Therefore we have that  $0 < d(H, K) \leq \inf\{d(x_n, y_n) | n \geq m\}$ .

(4) $\implies$ (5): Suppose that  $x \notin K$ , where  $K$  is compact, and  $x \in \overline{B(K; 1/n)}$  for each  $n \in \mathbb{N}$  with respect to the semimetric  $d$  satisfying the condition of (4). Then there exists a sequence  $\{z_n\}$  such that

$$z_n \in B(K; 1/n) \cap \text{int}B(x; 1/n).$$

Hence  $\{z_n\} \rightarrow x$ . Also  $d(x_n, z_n) < 1/n$  for some sequence  $\{x_n\} \subset K$ . Then there exist subsequences  $\{x_{n(i)}\} \subset \{x_n\}$  and  $\{z_{n(i)}\} \subset \{z_n\}$  such that  $\{x_{n(i)}\} \rightarrow p$  for some  $p \in K$  and  $\{z_{n(i)}\} \rightarrow x$ . Therefore there exist

$j, m \in \mathbb{N}$  such that  $d(x_{n(i)}, z_{n(i)}) \geq 1/m$  for each  $i \geq j$ . On the other hand,  $d(x_{n(k)}, z_{n(k)}) < 1/n(k)$  for some  $n(k) \geq \max\{n(j), m\}$ , which is a contradiction.

(5) $\implies$ (1): Let  $d$  be a compatible semimetric satisfying the condition of (5). If  $H$  and  $K$  are disjoint compact subsets of  $X$  with  $d(H, K) = 0$ , then  $\lim d(x_n, y_n) = 0$  for some sequences  $\{x_n\} \subset H$ ,  $\{y_n\} \subset K$ . Since  $X$  is first countable, there exist subsequences  $\{x_{n(i)}\} \subset \{x_n\}$ ,  $\{y_{n(i)}\} \subset \{y_n\}$  and points  $x \in H$ ,  $y \in K$  satisfying  $\{x_{n(i)}\} \longrightarrow x$ ,  $\{y_{n(i)}\} \longrightarrow y$ . Since  $y \notin \overline{B(H; 1/m)}$  for some  $m \in \mathbb{N}$ , we have that  $B(y; 1/k) \cap B(H; 1/k) = \emptyset$  for some  $k \geq m$ . This contradicts the fact that  $d(x_{n(i)}, y_{n(i)}) < 1/k$  and  $d(y, y_{n(i)}) < 1/k$  for some  $n(i) \in \mathbb{N}$ .

**Remark 3.6.** (1) The space  $Y$  in Example 4.9 is  $c$ -stratifiable  $\beta$ , but not  $q$ , and the Sorgenfrey line is  $c$ -stratifiable  $q$ , but not  $\beta$ .

(2) The space  $X$  in Example 4.6 is Moore (hence,  $X$  has a  $G_\delta^*$ -diagonal), but not  $K$ -semimetrizable, and the Nagata space  $X$  in Example 4.9 is  $K$ -semimetrizable, but not Moore.

(3) The space  $Y$  in Example 4.9 is stratifiable (hence  $c$ -stratifiable) Fréchet as the perfect image of a Nagata space (hence,  $K$ -semimetrizable), but  $Y$  is not semimetrizable (not even  $q$ ).

**Proposition 3.7.** *Every  $K$ -semimetrizable space has a  $G_\delta^*$ -diagonal.*

*Proof.* By Theorems 2.3 and 3.5, let  $g$  be a  $cs$ -stratifiable  $q$ ,  $\beta$ -function of  $X$  such that  $\{g_n(x)\}$  is a neighbourhood base of  $x$ . For each  $n \in \mathbb{N}$ , we put  $\mathcal{G}_n = \{g_n(x) \mid x \in X\}$ . To see that the sequence  $\{\mathcal{G}_n\}$  is a  $G_\delta^*$ -diagonal, suppose that  $x \neq y \in \bigcap_{n \geq 1} \overline{st(x, \mathcal{G}_n)}$ . Then there exist  $z_n \in g_n(y) \cap st(x, \mathcal{G}_n)$  ( $n \geq 1$ ). Hence  $\{z_n\} \longrightarrow y$  and  $x, z_n \in g_n(x_n)$  for some sequence  $\{x_n\}$ . Then  $\{x_n\} \longrightarrow x$  and  $y \notin C = \{x_n \mid n \geq m\} \cup \{x\}$  for some  $m \in \mathbb{N}$ . Hence  $y \notin \overline{g_k(C)}$  for some  $k \geq m$ . This is a contradiction.

The following theorem gives a condition for strong  $\alpha$ -ness and  $c$ -stratifiability to be equivalent, and follows directly from Theorems 2.4, 2.8 and 3.5 and Proposition 3.7.

**Theorem 3.8.** *For an orthocompact  $\beta$ ,  $q$ -space, the following conditions are equivalent.*

- (1)  $X$  is  $K$ -semimetrizable.
- (2)  $X$  has a  $G_\delta^*$ -diagonal.
- (3)  $X$  is strongly  $\alpha$ .

(4)  $X$  is  $cs$ -stratifiable.

An analogue to Theorem 3.5 for the class of regular spaces follows directly from Theorem 2.3.

**Theorem 3.9.** *For a regular space  $X$ ,  $X$  is  $K$ -semimetrizable if, and only if, it is weak  $c$ -stratifiable  $q, \beta$ .*

We next give some partial answers to the question of Burke [6; Question 2] on what minimal topological condition on a Moore space (or semimetric space) will ensure that the space is  $K$ -semimetrizable.

**Theorem 3.10.** (1) *Every  $T_2$ , orthocompact developable space  $X$  is  $K$ -semimetrizable.*

(2) *Every regular orthocompact semistratifiable  $q$ -space (hence, regular orthocompact semimetrizable space)  $X$  is  $K$ -semimetrizable.*

(3) *Every regular orthocompact  $c$ -semistratifiable  $q, \beta$ -space  $X$  is  $K$ -semimetrizable.*

(4) *Every regular  $k$ -semistratifiable  $q$ -space  $X$  is  $K$ -semimetrizable.*

*Proof.* Since a developable  $T_2$ -space has a  $G_\delta(2)$ -diagonal, (1) follows from Theorems 2.8 and 3.5. Since every semistratifiable  $T_2$ -space has a  $G_\delta$ -diagonal, (2) follows from Theorems 2.8 and 3.5. For (3), since  $X$  is semistratifiable, (3) follows from (2). (4) follows from Theorems 2.3, 2.4 and 3.5.

**Remark 3.11.** (1) With regards to (2) of Theorem 3.10, it is already known [1; page 133] or [22; page 441], that every paracompact semimetrizable space is  $K$ -semimetrizable.

(2) In (2) and (3) of Theorem 3.10, we can not change orthocompactness to subparacompactness by Example 4.6.

(3) In (4) of Theorem 3.10, we already know that a space is regular  $k$ -semistratifiable  $q$  if, and only if, it is Nagata [31; Theorem 2.1]. But, we do not know whether every  $T_2$ ,  $k$ -semistratifiable  $q$ -space is  $c$ -stratifiable. (If this answer is affirmative, then every  $T_2$ ,  $k$ -semistratifiable  $q$ -space is first countable and Nagata.) The converse of (4) does not hold, because the space  $\Psi$  in Example 4.5 is not  $k$ -semistratifiable.

In the following theorem, the equivalence of (1) and (4) is proved in [22; Theorem 5.4].

**Theorem 3.12.** For a space  $X$ , conditions (1)-(5) are all equivalent and (5)  $\implies$  (6) holds.

(1)  $X$  is a full  $K$ -semimetrizable space.

(2)  $X$  has a development  $\{\mathcal{G}_n\}$  such that if  $K_1$  and  $K_2$  are disjoint compact subsets, then  $st(K_1, \mathcal{G}_m) \cap K_2 = \emptyset$  for some  $m \in \mathbb{N}$ .

(3)  $X$  has a development  $\{\mathcal{G}_n\}$  such that if  $p \notin C$ , where  $C$  is the union of a convergent sequence and any one point of its limit points, then  $p \notin \overline{st(C, \mathcal{G}_m)}$  for some  $m \in \mathbb{N}$ .

(4)  $X$  satisfies one of the equivalent conditions in Theorem 3.3.

(5)  $X$  is a  $w\theta$ ,  $\beta$ -space with a regular  $G_\delta$ -diagonal.

(6)  $X$  is a developable  $c$ -stratifiable space.

*Proof.* (1) $\implies$ (2): Let  $d$  be a compatible full  $K$ -semimetric on  $X$ . For each  $n \in \mathbb{N}$ , we put  $\mathcal{G}_n = \{\text{int}B(x; \epsilon) \mid \delta B(x; \epsilon) < 1/n\}$ . Then  $\{\mathcal{G}_n\}$  is a development of  $X$  since  $d$  is a full semimetric. For, suppose that  $x \in X$  and  $x_n \in st(x, \mathcal{G}_n) \setminus U$  ( $n \geq 1$ ) for some open neighbourhood  $U$  of  $x$ . Then  $x, x_n \in G_n$  and  $\delta G_n < 1/n$  for some  $G_n \in \mathcal{G}_n$ , which is a contradiction. Now, suppose that  $K_1$  and  $K_2$  are disjoint compact subsets and  $x_n \in st(K_1, \mathcal{G}_n) \cap K_2$  for each  $n \in \mathbb{N}$ . Then  $y_n \in G_n \cap K_1$  and  $x_n \in G_n$  for some  $G_n \in \mathcal{G}_n$ . Since  $\delta G_n < 1/n$  ( $n \geq 1$ ),  $\lim_{n \rightarrow \infty} d(x_n, y_n) = 0$ . This contradicts  $d(K_1, K_2) > 0$ .

(2) $\implies$ (3): Let  $\{\mathcal{G}_n\}$  be a development of  $X$  satisfying (2). To see that  $X$  is  $T_2$ . let  $x \neq y$  and  $x_n \in st(x, \mathcal{G}_n) \cap st(y, \mathcal{G}_n)$  for each  $n \in \mathbb{N}$ . Then  $\{x_n\} \rightarrow x$  and  $\{x_n\} \rightarrow y$ . Given any open neighbourhood  $U$  of  $x$  with  $y \notin U$ ,  $S = \{x_n \mid n \geq m\} \cup \{x\} \subset U$  for some  $m \in \mathbb{N}$ . Then  $st(y, \mathcal{G}_k) \cap S = \emptyset$  for some  $k \geq m$ . This contradicts  $\overline{\{x_n\}} \rightarrow y$ . Next, suppose that  $p \notin K$ , where  $K$  is compact, and  $p \in \bigcap_{n \geq 1} st(K, \mathcal{G}_n)$ . Then  $a_n \in st(p, \mathcal{G}_n) \cap st(K, \mathcal{G}_n)$  ( $n \geq 1$ ). Hence  $a_n \in st(x_n, \mathcal{G}_n)$  for some sequence  $\{x_n\}$  in a sequentially compact  $K$ , and  $\{x_n\}$  contains a subsequence  $\{x_{n(i)}\}$  converging to some point  $x \in K$ . Since  $X$  is  $T_2$ ,  $L = \{x_{n(i)} \mid n(i) \geq m\} \cup \{x\}$  and  $H = \{a_{n(i)} \mid n(i) \geq m\} \cup \{p\}$  are disjoint for some  $m \in \mathbb{N}$ . Therefore,  $a_{n(k)} \in st(L, \mathcal{G}_{n(k)}) \cap H = \emptyset$  for some  $n(k) \geq m$ , which leads to a contradiction.

(3) $\implies$ (4): Let  $\{\mathcal{G}_n\}$  be a development of  $X$  such that  $\mathcal{G}_{n+1}$  is a refinement of  $\mathcal{G}_n$  and satisfies (3). We now show that  $\{\mathcal{G}_n\}$  satisfies the 3-link property. Suppose that  $x \neq y$  and for each  $n \in \mathbb{N}$ , there exists  $G_n \in \mathcal{G}_n$  such that  $x_n \in G_n \cap st(x, \mathcal{G}_n)$  and  $y_n \in G_n \cap st(y, \mathcal{G}_n)$ . Since  $\{x_n\} \rightarrow x$ ,  $\{y_n\} \rightarrow y$  and  $X$  is  $T_2$ ,  $y \notin C = \{x_n \mid n \geq m\} \cup \{x\}$  for some  $m \in \mathbb{N}$ . Hence  $y \notin \overline{st(C, \mathcal{G}_k)}$  for some  $k \geq m$ . Then  $y_l \in X \setminus \overline{st(C, \mathcal{G}_k)}$  for some  $l \geq k$  and  $x_l \in C$ . Therefore,  $y_l \in G_l \subset st(x_l, \mathcal{G}_l) \subset st(C, \mathcal{G}_k)$ , which is a contradiction.

(4) $\implies$ (5): Let  $X$  be a  $w\Delta$ -space with a regular  $G_\delta$ -diagonal. Then  $X$  satisfies condition (5).

(5) $\implies$ (1): By Theorem 3.4,  $X$  is a developable space with a regular  $G_\delta$ -diagonal. Hence there exists a compatible semimetric  $d$  on  $X$  satisfying (I) and (II) of (3) in Theorem 3.3. Then  $d$  is full by (3) $\implies$ (1) of [32; Theorem 2]. To see that  $d$  is  $K$ -semimetric, suppose that  $d(K, H) = 0$  for some disjoint compact subsets  $K$  and  $H$ . Then there are sequences  $\{x_n\} \subset K$  and  $\{y_n\} \subset H$  such that  $\lim_{n \rightarrow \infty} d(x_n, y_n) = 0$ . On the other hand, since  $X$  is a  $q$ -space with a  $G_\delta^*$ -diagonal,  $X$  is first countable. Hence  $\{x_n\}$  ( $\{y_n\}$ ) contains a subsequence  $\{x_{n(i)}\}$  ( $\{y_{n(i)}\}$ ) converging to a point  $x \in K$  ( $y \in H$ , respectively). Hence there are  $k, m \in \mathbb{N}$  such that  $d(x_{n(i)}, y_{n(i)}) \geq 1/m$  for each  $i \geq k$  by (II). This is a contradiction. Finally, (5) $\implies$ (6) follows from Theorems 2.8 and 3.4.

**Remark 3.13.** (1) The space  $\Psi$  in Example 4.5 is Moore and  $K$ -semi-metrizable, but not full  $K$ -semimetrizable.

(2) Every  $w\Delta$ -space is  $w\theta$  and  $\beta$ . Although the converse is an open problem [18; Problem 4.10], (4)  $\iff$  (5) of Theorem 3.12 (or (2) of Theorem 3.4) may be a slight progress to [32; Theorem 2] ([17; Theorem 2.5], respectively).

(3) The space  $X$  in Example 4.8 is  $T_2$  metacompact, full  $K$ -semi-metrizable, but not regular.

**Question 3.14.** Is every normal metacompact, full  $K$ -semimetrizable space, metrizable?

We next investigate conditions for spaces to be developable and  $K$ -semi-metrizable.

**Theorem 3.15.** *Consider the following conditions for a space  $X$ .*

- (1)  $X$  is developable and  $K$ -semimetrizable.
- (2)  $X$  is  $K$ -semimetrizable  $w\theta$ .
- (3)  $X$  is  $cs$ -stratifiable  $w\theta$  and  $\beta$ .
- (4)  $X$  is strongly  $\alpha$ ,  $w\theta$  and  $\beta$ .
- (5)  $X$  is developable  $T_2$ .

*Then, (1), (2) and (3) are equivalent.*

*Moreover, if  $X$  is orthocompact, then all conditions are equivalent.*

*Proof:* (1) $\implies$ (2)  $\implies$ (3) are evident. For (3) $\implies$ (1),  $X$  is  $K$ -semimetrizable by Theorem 3.5. Since  $X$  is semistratifiable and  $\theta$  by Theorem 2.10,  $X$  is developable [18; Remark 4.8]. (4) $\implies$ (3) follows from Theorem 2.4, and (3) $\implies$ (5) is evident. Moreover, if  $X$  is orthocompact, (5) $\implies$ (4) follows from Theorem 2.8.

Martin [26] showed that a locally connected rim-compact  $T_2$ -space  $X$  is  $K$ -semimetrizable if, and only if, it is developable  $\gamma$ .

**Definition 3.16.** A space  $X$  is said to be *rim-compact* if each point of  $X$  has a neighbourhood base consisting of open subsets with compact boundaries. A space  $X$  is *locally connected* if each point of  $X$  has a neighbourhood base consisting of connected open subsets.

We need the following lemma.

**Lemma 3.17.** (1) *Every locally connected rim-compact weak  $c$ -stratifiable (or,  $cs$ -stratifiable) space  $X$  is a  $c$ -stratifiable  $\gamma$ -space.*

(2) *Every pseudocompact Tychonoff weak  $c$ -stratifiable (or,  $cs$ -stratifiable) space  $X$  is a  $c$ -stratifiable  $\gamma$ -space.*

*Proof.* (1): First, let  $g$  be a weak  $c$ -stratifiable function of  $X$ . Then, we can assume that  $g_n(x)$  is connected for every  $x \in X$  and each  $n \in \mathbb{N}$ . To see that  $X$  is a  $\gamma$ -space, we use the same method given in the proof of [26; Theorem 4]. Suppose that  $K \subset W$ , where  $K$  is non-empty compact and  $W$  is open. Then there is an open subset  $G$  such that  $K \subset G \subset W$  and the boundary  $\partial G$  of  $G$  is compact. Since  $K \cap \partial G = \emptyset$ ,  $g_m(K) \cap \partial G = \emptyset$  for some  $m \in \mathbb{N}$ . Let  $K = \cup\{K_\alpha | \alpha \in A\}$ , where  $K_\alpha$  is a connected component of  $K$ . Since  $g_m(K_\alpha)$  is connected for each  $\alpha \in A$ ,  $g_m(K) = \cup_{\alpha \in A} g_m(K_\alpha) \subset G$ . Hence  $g$  is a  $\gamma$ -function by [23; Theorem 2.1]. Since  $X$  is first countable,  $g$  is a  $c$ -stratifiable function by [22; Theorem 1.3]. Next, let  $g$  be a  $cs$ -stratifiable function of  $X$ . To see that  $\{g_n(x)\}$  is a neighbourhood base of  $x$  for every  $x \in X$ , in the above proof, let  $K$  be a single point  $x$ . Since  $\{x\} \cap \partial G = \emptyset$  and  $\partial G$  is compact, we have that  $\overline{g_m(x)} \cap \partial G = \emptyset$  for some  $m \in \mathbb{N}$ . This asserts that  $\overline{g_m(x)} \subset G$ , which implies that  $X$  is first countable and regular. Therefore  $X$  is  $c$ -stratifiable by Theorem 2.3, and hence  $X$  is a  $\gamma$ -space.

(2): Let  $g$  be a weak  $c$ -stratifiable function or a  $cs$ -stratifiable function of  $X$ . By regularity of  $X$ , we assume that  $\overline{g_{n+1}(x)} \subset g_n(x)$ . Since  $\bigcap_{n \geq 1} \overline{g_n(x)} = \{x\}$ ,  $X$  is first countable by [27; Lemma 2.3]. Hence  $X$  is  $c$ -stratifiable by Theorem 2.3 and hence,  $X$  is  $\gamma$  by [22; Theorem 4.2].

**Theorem 3.18.** *Let  $X$  be a locally connected rim-compact space or a pseudocompact Tychonoff space. Then the following conditions are equivalent.*

- (1)  $X$  is developable and  $K$ -semimetrizable.
- (2)  $X$  is  $K$ -semimetrizable.
- (3)  $X$  is  $T_2$ , developable and  $\gamma$ .
- (4)  $X$  is weak  $c$ -stratifiable and  $\beta$ .



- (5)  $X$  is  $cs$ -stratifiable and  $\beta$ .
- (6)  $X$  is  $T_2$ ,  $\gamma$  and  $\beta$ .

*Proof.* First, we note that every  $\gamma, \beta$ -space is developable [18; Proposition 4.2]. (1) $\iff$ (4) and (1) $\iff$ (5) follow from Theorem 3.5 and Lemma 3.17.

(1) $\implies$ (2) $\implies$ (4) and (1) $\implies$ (3) $\implies$ (6) is evident. Since every  $T_2, \gamma$ -space is  $c$ -stratifiable, (6) $\implies$ (5) is true.

By the proof of the above theorem and Theorem 3.5, we have that in the class of  $T_2, \gamma$ -spaces, the following properties are coincident: (1) developable and  $K$ -semimetrizable, (2)  $K$ -semimetrizable, (3) developable and (4)  $\beta$ .

The next theorem follows from Theorem 3.10.

**Theorem 3.19.** *For an orthocompact  $T_2$ -space  $X$ ,  $X$  is developable and  $K$ -semimetrizable if, and only if, it is developable*

A Tychonoff space  $X$  is called a  $p$ -space [2] if in the Stone-Ćech compactification  $\beta X$ , there is a sequence  $\{\mathcal{G}_n\}$  of open covers of  $X$  such that  $\bigcap_{n \geq 1} st(x, \mathcal{G}_n) \subset X$  for every  $x \in X$ . Every locally compact  $T_2$ -space is a  $p$ -space.

Burke [5] showed that there is a locally compact  $T_2$ -space with a  $G_\delta$ -diagonal, which is not  $w\Delta$ . But, it is known that every locally compact semistratifiable  $T_2$ -space or every  $\theta$ -refinable  $p$ -space with a  $G_\delta$ -diagonal is Moore [8, 21]. Then we have the following result by Theorem 3.10.

**Theorem 3.20.** *For a metacompact  $p$ -space  $X$ ,  $X$  is Moore and  $K$ -semi-metrizable if, and only if, it has a  $G_\delta$ -diagonal.*

The next result was studied by Kotake [20] in the class of regular spaces.

**Theorem 3.21.** *For a space  $X$ , the following conditions are equivalent.*

- (1)  $X$  is Nagata.
- (2)  $X$  is  $K$ -semimetrizable  $wcc$ .
- (3)  $X$  is  $cs$ -stratifiable  $wN$ .
- (4)  $X$  is strongly  $\alpha$ ,  $wN$ .
- (5)  $X$  is a  $wN$ -space with a  $G_\delta^*$ -diagonal.
- (6)  $X$  is regular semimetrizable  $wcc$ .

*Proof.* Every Nagata space is stratifiable and first countable, hence it is  $c$ -stratifiable  $q$  and  $\beta$ . Therefore (1) $\implies$ (2) and (2) $\implies$ (3) follow from

Proposition 1.4 and Theorem 3.5, and (3) $\implies$ (1) follows from Theorem 2.10. (1) $\implies$ (4) and (4) $\implies$ (3) follow from Theorems 2.4 and 2.8. Also, (1) $\implies$ (5) is evident. To prove (5) $\implies$ (4), let  $g$  be a  $wN$ -function and  $\{\mathcal{G}_n\}$  be a  $G_\delta^*$ -diagonal sequence. Since regularity is not used to show that every  $\beta$ -space with a  $G_\delta^*$ -diagonal is semistratifiable [17; Theorem 5.2],  $X$  is a subparacompact  $wN$ -space. Then  $X$  is metacompact by [18; Corollary 3.5]. Hence  $X$  is strongly  $\alpha$  by Theorem 2.8. (1) $\implies$ (6) is evident. Finally, since every regular semistratifiable space has a  $G_\delta^*$ -diagonal [14; Theorem 5.11], (6) $\implies$ (5) follows from Proposition 1.4.

Regarding Question 2.12, we have the following corollary which follows from the fact that every  $wcc$ -space is  $\beta$ .

**Corollary 3.22.** *For a  $wN$ -space, the classes of the following spaces are all coincident.*

(1) Nagata spaces, (2) strongly  $\alpha$ -spaces, (3)  $c$ -stratifiable spaces, (4)  $K$ -semimetrizable spaces and (5) spaces with a  $G_\delta^*$ -diagonal.

**Remark 3.23.** Ceder [7; page 114] asked whether every paracompact semimetrizable space must be a Nagata space. Heath [16] showed that there is a paracompact  $K$ -semimetrizable cosmic (the continuous image of a separable metric space) space which is not a stratifiable space (hence, neither  $k$ -semistratifiable [24; Example 4.2] nor  $wcc$ ). He also posed the following problem: What topological condition is necessary for a paracompact semimetrizable (=  $K$ -semimetrizable) space to be an  $M_3$ -space? As a remark to this problem, one can note that in the class of regular semimetrizable spaces, Nagata spaces,  $k$ -semistratifiable spaces and  $wcc$ -spaces are coincident.

#### 4. METRIZABILITIES AND EXAMPLES

We begin this section with metrizations of  $wM$ -spaces. The concept of  $wM$ -spaces was given by Ishii [19]. Here we define a  $wM$ -space by an equivalent condition given by Hodel.

**Definition 4.1** [18; Theorem 5.2]. A space  $X$  is  $wM$  if, and only if, it is  $w\gamma$  and  $wN$ .

The following implications are well known.

An  $M$ -space (in the sense of Morita)  $\implies$  a  $wM$ -space  $\implies$  a  $w\Delta$ -space.

The class of  $wM$ -spaces is contained in the class of  $w\theta$ ,  $wcc$ -spaces. Therefore, we consider metrizations for the class of  $w\theta$ ,  $wcc$ -spaces. Metrizations

for this class was studied in [28]. For metrizations of  $wM$ -spaces, Martin [25] proved that every regular  $c$ -semistratifiable  $wM$ -space is metrizable, and Ishii [19] proved that every normal  $wM$ -space with a  $G_\delta^*$ -diagonal is metrizable. On the other hand, the space  $\Psi$  in Example 4.4 is a  $c$ -stratifiable Moore  $\gamma$ -space which is not metrizable.

**Theorem 4.2.** *Let  $X$  be a  $w\theta$ ,  $wcc$ -space. Then  $X$  is metrizable if  $X$  satisfies any one of the following statements.*

- (1)  $X$  is  $K$ -semimetrizable.
- (2)  $X$  is strongly  $\alpha$ .
- (3)  $X$  is  $cs$ -stratifiable.
- (4)  $X$  has a  $G_\delta^*$ -diagonal.
- (5)  $X$  is regular  $c$ -semistratifiable.

*Proof.* For all conditions (1)-(5),  $X$  is a  $wN$ -space by Proposition 1.4. Hence for (1)-(4),  $X$  is a  $w\theta$ , Nagata space by Theorem 3.21. Therefore,  $X$  is metrizable [30; Theorem 5]. For (5), since every  $wcc$ -space is  $\beta$ ,  $X$  is regular  $c$ -semistratifiable  $\beta$ , hence  $X$  is semistratifiable. Then  $X$  is  $wcc$  Moore [18; Corollary 4.6], which implies that  $X$  is metrizable [31; Corollary 3.6].

**Remark 4.3.** In Theorem 4.2, the condition  $w\theta$  ( $wcc$ ) can not be weakened to  $q$  ( $\beta$ , respectively). Indeed, the Nagata-space  $X$  in Example 4.9 is a  $q$ ,  $wcc$ -space which satisfies all of the conditions (1)-(5) in Theorem 4.2, but is not metrizable. Also, the space  $\Psi$  in Example 4.5 is a  $\gamma$ ,  $\beta$ -space which satisfies all of the conditions (1)-(5) in Theorem 4.2, but is not metrizable.

The second part (2) of the next theorem is a generalization of Lee's result [22] that every pseudocompact Tychonoff stratifiable space is metrizable.

**Theorem 4.4.** (1) *Every locally connected rim-compact  $k$ -semistratifiable space  $X$  is metrizable.*

(2) *Every pseudocompact Tychonoff  $k$ -semistratifiable space  $X$  is metrizable.*

*Proof.* First, we show that if  $X$  satisfies the conditions of (1), then  $X$  is a first countable  $T_2$ -space. Let  $g$  be a  $k$ -semistratifiable function such that  $g_n(x)$  is connected. To see that  $\{g_n(x)\}$  is a neighbourhood base of  $x$  for every  $x \in X$ , suppose that  $x \in U$  and  $g_n(x) \setminus U \neq \emptyset$  ( $n \geq 1$ ), where  $U$  is open. Then there is an open neighbourhood  $W$  of  $x$  such that  $W \subset U$  and the boundary  $\partial W$  is compact. Since  $g_m(x) \cap \partial W = \emptyset$  for some  $m \in \mathbb{N}$ ,

$g_m(x) = (g_m(x) \cap W) \cup (g_m(x) \setminus \overline{W})$  is not connected. This contradiction implies that  $\{g_n(x)\}$  is a neighbourhood base of  $x$ . To see that  $X$  is Hausdorff, let  $x \neq y$  and  $x_n \in g_n(x) \cap g_n(y)$  ( $n \geq 1$ ). Then for any open neighbourhood  $U$  of  $x$  with  $y \notin U$ ,  $K = \{x_n | n \geq m\} \cup \{x\} \subset U$  for some  $m \in \mathbb{N}$ . Hence  $g_l(y) \cap K = \emptyset$  for some  $l \geq m$ , which is a contradiction. Next, in both cases,  $X$  is a  $\gamma$ -space by Theorem 2.4 and Lemma 3.17. Also,  $X$  is a *wcc*-space. Indeed, let  $g$  be a  $k$ -semistratifiable function such that, whenever  $b_n \in g_n(a_n)$  ( $n \geq 1$ ) and  $\{b_n\} \rightarrow b$ , then  $\{a_n\} \rightarrow b$ . Now, suppose that  $y_n \in g_n(x_n)$  ( $n \geq 1$ ) and  $\{y_n\}$  has a cluster point  $y$ . Since  $X$  is first countable, there exists a subsequence  $\{y_{n(i)}\}$  of  $\{y_n\}$  converging to  $y$  and  $y_{n(i)} \in g_i(x_{n(i)})$  ( $n \geq 1$ ). Hence  $\{x_{n(i)}\}$  converges to  $y$ , which implies that  $g$  is a *wcc*-function. Finally, every  $\gamma$ , *wcc*  $T_2$ -space is metrizable [31; Corollary 3.6].

We note that Martin [26; Example 3] showed that there exists a locally connected locally compact  $K$ -semimetrizable Moore space  $X$  which is not normal. This space is not *wcc* by Theorem 3.21.

As regards to Theorem 4.4, (2) is proved in [30; Corollary 4] in a different way, and as for (1), every locally compact  $T_2$  (even sieve-complete regular)  $k$ -semistratifiable is metrizable [30; Theorem 18].

**Example 4.5.** [22; Example 6.6] The space  $\Psi$  in [13; 51] is Moore and  $K$ -semimetrizable that is not full  $K$ -semimetrizable. First, it is known that  $\Psi$  is a locally compact pseudocompact separable Moore  $c$ -stratifiable space that is not metacompact. To see that  $\Psi$  is orthocompact, for any  $E = \{x_k^E | k \in \mathbb{N}\} \in \mathcal{E}$ , where  $\{x_k^E | k \in \mathbb{N}\}$  is an infinite subsequence of  $\mathbb{N}$ , we put  $B(\omega_E, n) = \{\omega_E\} \cup \{x_n^E, x_{n+1}^E, \dots\}$  ( $n \in \mathbb{N}$ ). Then any open cover  $\mathcal{G}$  of  $\Psi$  has the refinement  $\mathcal{H} = \{\{n\} | n \in \mathbb{N}\} \cup \{B(\omega_E, n(E)) | E \in \mathcal{E}\}$ , where for any  $E \in \mathcal{E}$ ,  $B(\omega_E, n(E)) \subset G$  for some  $G \in \mathcal{G}$  and some  $n(E) \in \mathbb{N}$ . And  $\cap \mathcal{W}$  is open for any  $\mathcal{W} \subset \mathcal{H}$ . Therefore,  $\Psi$  is strongly  $\alpha$  by Theorem 2.8. Then  $\Psi$  is  $K$ -semimetrizable and  $\gamma$  by Theorem 3.5 and Lemma 3.17. But  $\Psi$  does not have a regular  $G_\delta$ -diagonal [27; Theorem 2.6], and not *wcc* from Theorem 3.21. Hence it is not full  $K$ -semimetrizable by Theorem 3.12 and not  $k$ -semistratifiable since every first countable  $k$ -semistratifiable space is *wcc*.

**Example 4.6.** [6] Burke constructed the separable Moore (hence, semimetrizable) space  $X$  which is not  $K$ -semimetrizable. Hence,  $X$  is a  $c$ -semistratifiable  $\alpha$ -space which is neither strongly  $\alpha$  nor  $cs$ -stratifiable by Theorems 2.4 and 3.5. Also,  $X$  is not metacompact by Theorem 2.8 and not

$\gamma$ .

**Example 4.7.** [18; Example 4.14]. The Sorgenfrey line  $K$  is a paracompact  $\gamma$ -space with a  $G_\delta$ -diagonal. Hence  $K$  is strongly  $\alpha$  and  $c$ -stratifiable, but not semistratifiable (not even  $\beta$  [18; Proposition 4.2]).

**Example 4.8.** [9: Example 5.3.4] There exists a metacompact full  $K$ -semimetrizable space which is neither  $wcc$  nor regular.

Indeed, let  $X$  be the space of real numbers with the topology generated by the neighbourhood system  $\{\mathcal{U}(x)|x \in X\}$ , where  $\mathcal{U}(x) = \{U_n(x)|n \in \mathbb{N}\}$  and

$$U_n(x) = \begin{cases} (x - 1/n, x + 1/n) & \text{if } x \neq 0, \\ (x - 1/n, x + 1/n) \setminus \{1/k|k \in \mathbb{Z} \setminus \{0\}\} & \text{if } x = 0, \end{cases}$$

where  $\mathbb{Z}$  denotes the set of integers. It is well known that  $X$  is a metacompact  $T_2$ -space which is not regular. For each  $x \in X$ , we put

$$W_n(x) = \begin{cases} U_n(x) \setminus \{0\} & \text{if } x \neq 0, \\ U_n(x) & \text{if } x = 0. \end{cases}$$

Let  $\mathcal{W}_n = \{W_n(x)|x \in X\}$  for each  $n \in \mathbb{N}$ . Then it is easily seen that the sequence  $\{\mathcal{W}_n\}$  is a development satisfying the 3-link property. Therefore,  $X$  is full  $K$ -semimetrizable. Then  $X$  is strongly  $\alpha$  and  $c$ -stratifiable by Theorem 2.8. Also, if  $X$  is  $wcc$ , then it is metrizable by Theorem 4.2, which is a contradiction.

**Example 4.9.** [24; Example 4.3] There exist a first countable stratifiable space  $X$  and a perfect map  $f$  from  $X$  onto a non- $q$ -space  $Y$ . Then  $X$  is a Nagata space (hence,  $X$  is  $K$ -semimetrizable) which is not  $w\theta$  [30; Theorem 5] and  $Y$  is a stratifiable space which is not  $q$ . Then,  $Y$  is strongly  $\alpha$  and  $c$ -stratifiable but not semimetrizable.

**Example 4.10.** [10; Example 4.2] A regular full  $K$ -semimetrizable space that is not orthocompact. Let  $R = \{(x, y)|x, y \text{ are rational and } y > 0\}$ . Let  $J$  be the set of irrational numbers and let  $X = R \cup (J \times \{0\})$ . We give  $R$  the usual subspace topology  $\mathcal{T}^*$ . For each  $x \in J$  and each  $\epsilon > 0$ , let  $B(x, \epsilon) = \{(x, 0)\} \cup \{(x + k, h)||k| < h < \epsilon\}$ . Then  $\mathcal{T}^* \cup \{B(x, \epsilon)|x \in J, \epsilon > 0\}$  is a basis for a topology on  $X$ . Then  $X$  is a separable Moore space that is not orthocompact. Also,  $X$  has a development satisfying the 3-link property, hence full  $K$ -semimetrizable and  $c$ -stratifiable.

But I don't know whether this space is strongly  $\alpha$ .

## REFERENCES

- [1] Arhangel'skiĭ A. V, *Mappings and spaces*, Russian Math. Surveys, **21** (1966), 115-162.
- [2] Arhangel'skiĭ A. V, *On a class of spaces containing all metric spaces and all locally bicomact spaces*, Amer. Math. Soc. Transl., **92** (1970), 1-39.
- [3] Borges C. J. R, *On stratifiable spaces*, Pacific J. Math., **17** (1966), 1-16.
- [4] Borges C. J. R, *On metrizable spaces*, Canad. J. Math., **20** (1968), 795-804.
- [5] Burke D. K, *A nondevelopable locally compact Hausdorff space with a  $G_\delta$ -diagonal*, General Top. and its Appl., **2** (1972), 287-291.
- [6] Burke D. K, *On  $K$ -semimetric spaces*, Proc. Amer. Math. Soc., **73** (1979), 126-128.
- [7] Ceder. J. G, *Some generalizations of metric spaces*, Pacific J. Math., **11** (1961), 105-125.
- [8] Creede G. D, *Concerning semi-stratifiable spaces*, Pacific J. Math., **32** (1970), 47-54.
- [9] Engelking R, *General Topology*, Polish Sci. Publ., Warsaw, 1977.
- [10] Fletcher P and Lindgren W. F, *Orthocompactness and Strong Čech completeness in Moore spaces*, Duke Math. J., **39** (1972), 753-766.
- [11] Fletcher P and Lindgren W. F,  *$\theta$ -spaces*, General Topology and its Appl., **9** (1978), 139-153.
- [12] Gao Z, *On  $g$ -function separation*, Questions and Answers in Gen. Topology, **4** (1986), 47-56.
- [13] Gillman L and Jerison M, *Rings of continuous functions*, D. Van Nostrand, Princeton, N.J., 1960.
- [14] Gruenhage G, *Generalized metric spaces*, Chap. 10, in: K. Kunen and J. E. Vaughan, eds., Handbook of set theoretic topology, (North-Holland, Amsterdam).
- [15] Heath R. W, *On open mappings and certain spaces satisfying the first countability axiom*, Fund. Math., **57** (1965), 91-96.
- [16] Heath R. W, *A paracompact semimetric space which is not an  $M_3$ -space*, Proc. Amer. Math. Soc., **17** (1966), 868-870.
- [17] Hodel R. E, *Moore spaces and  $w\Delta$ -spaces*, Pacific J. Math., **38** (1971), 641-652.
- [18] Hodel R. E, *Spaces defined by sequences of open covers which guarantee that certain sequences have cluster points*, Duke Math. J., **39** (1972), 253-263.
- [19] Ishii T, *On  $wM$ -spaces I,II*, Proc. Japan Acad., **46** (1970), 5-10, 11-15.
- [20] Kotake Y, *On Nagata spaces and  $wN$ -spaces*, Sci. Rep. Tokyo Kyoiku Daigaku, Sec. A, **12** (1973), 46-48.
- [21] Kullman D. E, *Developable spaces and  $p$ -spaces*, Proc. Amer. Math. Soc., **27** (1971), 154-160.
- [22] Lee K. B, *Spaces in which compacta are uniformly regular  $G_\delta$* , Pacific J. Math., **81** (1979), 435-446.
- [23] Lindgren W. F and Fletcher P, *Locally quasi-uniform spaces with countable bases*, Duke Math. J., **41** (1974), 231-240.
- [24] Lutzer J, *Semistratifiable and stratifiable spaces*, General Top. and its Appl., **1** (1971), 43-48.
- [25] Martin H. W, *Metrizability of  $M$ -spaces*, Canad. J. Math., **25** (1973), 840-841.
- [26] Martin H. W, *Local connectedness in developable spaces*, Pacific J. Math., **61** (1975), 219-224.
- [27] McArthur W. G,  *$G_\delta$ -diagonals and metrization theorems*, Pacific J. Math., **44** (1973), 613-617.

- [28] Nitta S and Yoshioka I, *Nagata spaces and  $wN$ -spaces which are preserved by quasi-perfect maps*, Houston J. Math., **32** (2006), 1061-1075.
- [29] Okuyama A, *Some generalizations of metric spaces, their metrization theorems and product spaces*, Sci. Rep. Tokyo Kyoiku Daigaku Sect. A, **9** (1967), 236-254.
- [30] Yoshioka I, *On the metrizations of  $\gamma$ -spaces and  $ks$ -spaces*, Questions and Answers in Gen. Topology, **19** (2001), 55-72.
- [31] Yoshioka I, *Closed images of spaces having  $g$ -functions*, Topology and its Appl., to appear.
- [32] Zenor P, *On spaces with regular  $G_\delta$ -diagonals*, Pacific J. Math., **40** (1972), 759-763.

IWAO YOSHIOKA

139 MITSUYOSHI, KOURYO-CHO, KITA-KATSURAGI, NARA 635-0823, JAPAN

*e-mail address:* yoshioka.iwao@ybb.ne.jp

*(Received March 6, 2007)*

*(Revised June 5, 2007)*