

## A REMARK ON THE LAVALLEE-SPEARMAN-WILLIAMS-YANG FAMILY OF QUADRATIC FIELDS

KWANG-SEOB KIM AND YASUHIRO KISHI

ABSTRACT. In [4], M. J. Lavallee, B. K. Spearman, K. S. Williams and Q. Yang introduced a certain parametric  $D_5$ -quintic polynomial and studied its splitting field. The present paper gives an infinite family of quadratic fields with class number divisible by 5 by using properties of its polynomial.

### 1. INTRODUCTION

One of the main topics in algebraic number theory is to study the structure of ideal class groups of number fields, particularly the divisibility of class numbers. In the case of quadratic fields, numerous related results are known by many authors. As was first shown by T. Nagell [7] (resp. by Y. Yamamoto [11]), there exist infinitely many imaginary (resp. real) quadratic fields with class numbers divisible by an arbitrary given integer  $n$ . After that, several authors gave such results (for example, [1], [10], [6], [9], [12], [5], and so on).

Every unramified cyclic extension of prime degree  $p$  of a quadratic field is normal over  $\mathbb{Q}$  and is a  $D_p$ -extension of  $\mathbb{Q}$ , that is, a Galois extension whose Galois group is isomorphic to the dihedral group  $D_p$  of order  $2p$ . Therefore, it is given by roots of a polynomial of degree  $p$  with rational coefficients. Y. Kishi and K. Miyake [2] used some  $D_3$ -cubic polynomials to classify all quadratic fields whose class number is divisible by 3. Here, a  $D_3$ -cubic polynomial means a polynomial of degree 3 such that its minimal splitting field over  $\mathbb{Q}$  is a  $D_3$ -extension of  $\mathbb{Q}$ . In the case  $p = 5$ , Kishi [3] found a family of  $D_5$ -quintic polynomials such that its minimal splitting fields over  $\mathbb{Q}$  are unramified cyclic quintic extensions of quadratic subfields. But his methods is slightly complicated. In this paper, we will give such polynomials that are simpler and more efficient by using M. J. Lavallee, B. K. Spearman, K. S. Williams and Q. Yang's results.

For a number field  $K$ , let  $D(K)$  denote the discriminant of  $K$ . For a polynomial  $f$ , let  $d(f)$  (resp.  $\text{Spl}_{\mathbb{Q}}(f)$ ) denote the discriminant (resp. the minimal splitting field over  $\mathbb{Q}$ ) of  $f$ .

---

*Mathematics Subject Classification.* Primary 11R29; Secondary 11R11.

*Key words and phrases.* Class numbers, Quadratic fields,  $D_5$ -polynomials.

## 2. MAIN THEOREM

We need the following well-known lemma before we state our main theorem.

**Lemma 1** (Dedekind). *Let  $F$  be an algebraic number field and  $\mathfrak{D}$  be the different of  $F$  over  $\mathbb{Q}$ . Let  $\mathfrak{p}$  be a prime divisor in  $F$  over a prime number  $p$ , and  $\mathfrak{p}^d \parallel \mathfrak{D}$  and  $\mathfrak{p}^e \parallel p$ . Then*

- (a) *if  $p \nmid e$ , then  $d = e - 1$ ,*
- (b) *if  $p^v \parallel e$  ( $v > 0$ ), then  $e \leq d \leq ev + e - 1$ .*

*Proof.* See, for example, [8, Chap. III, §2, (2.6) Theorem]. □

**Theorem.** *Let  $f$  be a  $D_5$ -quintic polynomial and put  $L := \text{Spl}_{\mathbb{Q}}(f)$ . Moreover, let  $K$  be the unique quadratic subfield of  $L$ . Suppose that*

$$5^6 \nmid d(f) \quad \text{and} \quad q^4 \nmid d(f)$$

*for every prime number  $q \neq 5$ . Then  $L$  is an unramified cyclic quintic extension of  $K$ . Especially, the class number of  $K$  is divisible by 5.*

*Proof.* Let  $F = \mathbb{Q}(\theta)$  where  $f(\theta) = 0$ . Then we have

$$d(f) = a^2 D(F)$$

for some integer  $a$ .

If  $L/K$  is unramified, then the class number of  $K$  is divisible by 5 by class field theory. So it suffices to show that if  $L/K$  is not unramified, then we have

$$5^6 \mid D(F) \quad \text{or} \quad q^4 \mid D(F)$$

for some prime number  $q \neq 5$ . Suppose that  $\mathfrak{p}$  is a prime ideal of  $K$  which is ramified in  $L/K$ .

Case 1, where  $5 \mid \mathfrak{p}$ . Since  $\mathfrak{p}$  is ramified in  $L/K$  and  $[L : K] = 5$ , 5 is wildly ramified in  $F/\mathbb{Q}$ . By Lemma 1 (b), it must hold that  $5 \leq d \leq 9$ . This implies that  $5^5 \mid D(F)$ . Now we may view  $\text{Gal}(L/K)$  as a subgroup of the symmetric group  $S_5$  of order 5 because  $f$  is a quintic polynomial. Since  $\text{Gal}(L/K) \simeq D_5$  is contained in the alternative group  $A_5$  of degree 5,  $D(F)$  must be a square in  $\mathbb{Z}$ . Thus we have  $5^6 \mid D(F)$ .

Case 2, where  $q \mid \mathfrak{p}$  with  $q \equiv 1 \pmod{5}$ . Since  $\mathfrak{p}$  is ramified in  $L/K$  and  $[L : K] = 5$ ,  $q$  is tamely ramified in  $F/\mathbb{Q}$ . By Lemma 1 (a), we have  $q^4 \parallel \mathfrak{D}$ . In this case, we can easily verify that  $q^4 \mid D(F)$ . The proof is completed. □

## 3. THE LAVALLEE-SPEARMAN-WILLIAMS-YANG FAMILY OF QUADRATIC FIELDS

By using our theorem, we will give an infinite family of quadratic fields with class number divisible by 5. This is based on Lavallee, Spearman, Williams and Yang's works [4]. Let us refer to them.

For an integer  $b$ , we define

$$f_b(x) := x^5 - 2x^4 + (b + 2)x^3 - (2b + 1)x^2 + bx + 1.$$

First they proved that  $f_b$  is irreducible for any  $b \in \mathbb{Z}$  and

$$d(f_b) = (4b^3 + 28b^2 + 24b + 47)^2$$

(see [4, Lemmas 2.1 and 2.2]). They also proved the following.

**Proposition 1** ([4, Lemmas 2.3 and 2.4]). *If  $-(4b^3 + 28b^2 + 24b + 47)$  is not a square in  $\mathbb{Z}$ , then  $\text{Spl}_{\mathbb{Q}}(f_b)$  is a  $D_5$ -extension of  $\mathbb{Q}$  and the unique quadratic subfield of  $\text{Spl}_{\mathbb{Q}}(f_b)$  is*

$$\mathbb{Q}(\sqrt{-(4b^3 + 28b^2 + 24b + 47)}).$$

Furthermore, there are infinitely many integers  $b$  such that  $-(4b^3 + 28b^2 + 24b + 47)$  is square-free, is not a square in  $\mathbb{Z}$ , and the quadratic fields  $\mathbb{Q}(\sqrt{-(4b^3 + 28b^2 + 24b + 47)})$  are distinct, as we can see in the proof of [4, Theorem]. From this, together with Proposition 1 and our main theorem, the following proposition is immediately obtained.

**Proposition 2.** *There exist infinitely many quadratic fields of the form*

$$\mathbb{Q}(\sqrt{-(4b^3 + 28b^2 + 24b + 47)}), \quad b \in \mathbb{Z}$$

*whose class number is divisible by 5.*

#### ACKNOWLEDGEMENT

The authors would like to thank the referee for his careful reading of the manuscript and for giving many valuable comments.

#### REFERENCES

- [1] N. Ankeny and S. Chowla, *On the divisibility of the class numbers of quadratic fields*, Pacific J. Math. **5** (1955), 321–324.
- [2] Y. Kishi and K. Miyake, *Parametrization of the quadratic fields whose class numbers are divisible by three*, J. Number Theory **80** (2000), no. 2, 209–217.
- [3] Y. Kishi, *A new family of imaginary quadratic fields whose class number is divisible by five*, J. Number Theory **128** (2008), no. 8, 2450–2458.
- [4] M. J. Lavalée, B. K. Spearman, K. S. Williams and Q. Yang, *Dihedral quintic fields with a power basis*, Math. J. Okayama Univ. **47** (2005), 75–79.
- [5] S. R. Louboutin, *On the divisibility of the class number of imaginary quadratic number fields*, Proc. Amer. Math. Soc. **137** (2009), no. 12, 4025–4028.
- [6] M. R. Murty, *Exponents of class groups of quadratic fields*, in: “Topics in Number Theory,” (University Park, PA, 1997), pp. 229–239, Mathematical Applications, **467**, Kluwer Acad. Publ., Dordrecht, 1999.
- [7] T. Nagell, *Über die Klassenzahl imaginär quadratischer Zahlkörper*, Abh. Math. Seminar Univ. Hambrug **1** (1922), no. 1, 140–150.

- [8] J. Neukirch, “Algebraic number theory,” Grundlehren der Mathematischen Wissenschaften, **322**, Springer-Verlag, Berlin, 1999.
- [9] K. Soundararajan, *Divisibility of class numbers of imaginary quadratic fields*, J. London Math. Soc. (2) **61** (2000), no. 3, 681–690.
- [10] P. Weinberger, *Real quadratic fields with class numbers divisible by  $n$* , J. Number Theory **5** (1973), 237–241.
- [11] Y. Yamamoto, *On unramified Galois extensions of quadratic number fields*, Osaka J. Math. **7** (1970), 57–76.
- [12] G. Yu, *A note on the divisibility of class numbers of real quadratic fields*, J. Number Theory **97** (2002), no. 1, 35–44.

KWANG-SEOB KIM  
SCHOOL OF MATHEMATICS  
KOREA INSTITUTE FOR ADVANCED STUDY  
SEOUL, 130-722 KOREA  
*e-mail address*: kwang12@kias.re.kr

YASUHIRO KISHI  
DEPARTMENT OF MATHEMATICS  
AICHI UNIVERSITY OF EDUCATION  
AICHI, 448-8542 JAPAN  
*e-mail address*: ykishi@aeu.ac.jp

(*Received April 21, 2016*)  
(*Accepted August 22, 2016*)