

# ON CONNECTED LOCALLY CONNECTED SPACES WITH CUT POINTS

Dedicated to Professor TAKESHI INAGAKI on his 60th birthday

IWAOKA YOSHIOKA

If  $M$  is a connected space and  $p$  is a point of  $M$  such that  $M-p$  is not connected, then  $p$  will be called a *cut point* of  $M$  [1]. If  $p$  is a cut point,  $M-p$  is the sum of two mutually separated sets  $M_1(p)$  and  $M_2(p)$ , then  $M_1(p)$  and  $M_2(p)$  will be called *sects* of  $M$  from  $p$  [2]. A point  $q$  is said to be separated from a point  $r$  by  $p$  if there exists a separation

$$M-p = M_1(p) \cup M_2(p), \text{ where } M_1(p) \ni q, M_2(p) \ni r.$$

If  $r$  is a point of  $M$ , then  $S(r, M)$  denotes the set of all points which are separated from  $r$  by at least one cut point of  $M$ , and  $T(r, M)$  the set of all cut points of  $M$  except the point  $r$ . The boundary of a point set  $A$  will be denoted by  $Bd A$ . In this note, all spaces are connected locally connected Hausdorff spaces with at least one cut point. The purpose of this note is to study the conditions under which the set  $S(r, M)$  is connected.

The author is grateful to Prof. K. Koseki for his kind advice.

**Lemma 1.** *If  $p$  is a point of a space  $M$ , then  $Bd S(p, M) \subset \overline{T(p, M)}$ .*

*Proof.* Let us assume the contrary and let  $x \in Bd S(p, M) - \overline{T(p, M)}$ . Then, since  $M$  is locally connected, there exist neighbourhoods  $U(x)$ ,  $V(x)$  of a point  $x$  such that

$$x \in V(x) \subset C \subset U(x) \text{ and } U(x) \cap T(p, M) = \emptyset,$$

where  $C$  is a connected set.

On the other hand, there is a point  $y \in V(x) \cap S(p, M)$ . Therefore, we can take a point  $q \in T(p, M)$  and have a separation

$$M-q = M_1(q) \cup M_2(q), \text{ where } M_1(q) \ni p, M_2(q) \ni y.$$

Since  $C \ni y$  and  $C \not\ni q$ ,  $x \in M_2(q) \subset S(p, M)$ . This contradiction completes the proof.

**Lemma 2.** *Let  $p$  be a point of a space  $M$ . If  $S(p, M)$  is connected and  $Bd S(p, M) \cap T(p, M) \neq \emptyset$ , then  $Bd S(p, M)$  is a single point.*

*Proof.* Let  $q$  be a point of the set  $Bd S(p, M) \cap T(p, M)$ . Then we have a separation

$$M - q = M_1(q) \cup M_2(q), \text{ where } p \in M_1(q), M_2(q) \subset S(p, M).$$

Since  $q \in Bd S(p, M)$ ,  $S(p, M) \subset M - q$ . Therefore, we have  $M_2(q) = S(p, M)$ , because  $M_2(q)$  is open and closed in  $M - q$ . This means that  $Bd S(p, M)$  is a single point.

**Theorem 3.** *For a point  $p$  of a space  $M$ , the following four conditions are equivalent :*

- (a) *If  $S(p, M)$  is non-vacuous, it is a connected set.*
- (b) *If  $M_2(q), M_2(q')$  are two sects of  $M$  from  $q, q'$  not containing  $p$ , where  $q, q' \in T(p, M)$ , then there exists a connected sect of  $M$ , not containing  $p$ , which contains  $M_2(q) \cup M_2(q')$ .*
- (c) *If  $M_2(q), M_2(q')$  are two connected sects of  $M$  from  $q, q'$  not containing  $p$ , where  $q, q' \in T(p, M)$  then there exists a connected sect of  $M$ , not containing  $p$ , which contains  $M_2(q) \cup M_2(q')$ .*
- (d)  *$Bd S(p, M)$  is a single point such that if  $Bd S(p, M) \neq p$ ,  $M$  is uniquely separated by  $Bd S(p, M)$ , and if  $Bd S(p, M) = p$ , there exists a component  $K$  of  $S(p, M)$  with  $\overline{T(p, M) - K} \ni Bd S(p, M)$ .*

*Proof.* (a) implies (b): Let  $M_2(q), M_2(q')$  be sects satisfying the assumption of (b) and put

$$\mathfrak{A} = \{N_2(t) \mid M_2(q) \subset N_2(t)\}, \quad \mathfrak{A}^* = \bigcup_{N_2(t) \in \mathfrak{A}} N_2(t)$$

where  $N_2(t)$  is any sect of  $M$  from  $t$  not containing  $p$  for each point  $t \in T(p, M)$ . Let us assume that  $x \in S(p, M) - \mathfrak{A}^*$ . Then, there exists a cut point  $u$  with a separation

$$M - u = H_1(u) \cup H_2(u), \text{ where } H_1(u) \ni p, H_2(u) \ni x.$$

Then  $H_2(u) \cap \mathfrak{A}^* = \emptyset$ . For, if  $H_2(u) \cap \mathfrak{A}^* \neq \emptyset$ , there exists  $K_2(t) \in \mathfrak{A}$  and  $K_2(t) \cap H_2(u) \neq \emptyset$ . If  $u = t$ , then  $H_2(u) \cup K_2(t) \in \mathfrak{A}$ . This implies  $u \neq t$ . Since  $H_1(u) \cap K_1(t) \ni p$ ,

$$H_2(u) \subset K_2(t) \text{ or } H_2(u) \supset K_2(t).$$

Hence  $x \in \mathfrak{U}^*$ . This implies  $H_2(u) \cap \mathfrak{U}^* = \emptyset$ . By the connectedness of  $S(p, M)$ ,  $S(p, M) = \mathfrak{U}^*$ , because  $\mathfrak{U}^*$  is open and closed in  $S(p, M)$ . Therefore, there exists a sect  $N_2(t) \in \mathfrak{U}$  such that  $N_2(t) \cap M_2(q') \neq \emptyset$ . Then

- (1)  $N_2(t) \cup M_2(q') \supset M_2(q) \cup M_2(q')$  if  $t = q'$ ,  
 (2)  $M_2(q') \supset M_2(q) \cup M_2(q')$  or  $N_2(t) \supset M_2(q) \cup M_2(q')$  if  $t \neq q'$ .

(1) and (2) imply that there exists a sect  $L_2(s) \in \mathfrak{U}$  with  $L_2(s) \supset M_2(q) \cup M_2(q')$ . If there exists a sect  $H_2(r) \in \mathfrak{U}$  with  $H_2(r) \ni s$ , we consider a component  $C$  of  $H_2(r)$  containing  $s$ . Then

$$C \supset L_2(s) \supset M_2(q) \cup M_2(q').$$

If there exists no such sect, then  $s \notin S(p, M)$ . Therefore, by the connectedness of  $S(p, M)$ ,

$$L_2(s) = S(p, M) \supset M_2(q) \cup M_2(q').$$

It is clear that (b) implies (c) and (c) implies (a).

(a) implies (d): Suppose that  $Bd S(p, M)$  contains distinct points  $t_1$  and  $t_2$ . By Lemma 2,

$$(3) \quad Bd S(p, M) \cap T(p, M) = \emptyset.$$

As  $M$  is a Hausdorff space, there exist neighbourhoods  $U_i \ni t_i (i=1, 2)$  with  $U_1 \cap U_2 = \emptyset$ . By the locally connectedness of  $M$ , there exist neighbourhoods  $V_i$  and connected sets  $C_i$  satisfying  $U_i \supset C_i \supset V_i \ni t_i (i=1, 2)$ . By Lemma 1, there exist  $q_i \in V_i \cap T(p, M) (i=1, 2)$ . Then  $q_i \in S(p, M) (i=1, 2)$  by  $T(p, M) \subset \overline{S(p, M)}$  and (3). Let  $H_2(u_i)$  be connected sects of  $M$  from  $u_i$  not containing  $p$  with  $u_i \in T(p, M)$  and  $q_i \in H_2(u_i) (i=1, 2)$ . Since (a) and (c) are equivalent, there exists a connected sect  $H_2(u_3)$  of  $M$  from  $u_3 \in T(p, M)$  not containing  $p$  such that  $H_2(u_1) \cup H_2(u_2) \subset H_2(u_3)$ . Hence

$$H_2(u_3) \cap C_1 \neq \emptyset \neq H_2(u_3) \cap C_2.$$

Then  $C_1 \cap C_2 \ni u_3$ , because  $C_1$  and  $C_2$  are connected and  $H_2(u_3)$  is open and closed in  $M - u_3$ . This implies that  $Bd S(p, M)$  is a single point. Now, let  $t = Bd S(p, M)$ . If  $t \neq p$ , there exists a separation such that

$$M - t = \{M - \overline{S(p, M)}\} \cup S(p, M), \text{ where } M - \overline{S(p, M)} \ni p.$$

This separation is uniquely determined. On the other hand, if  $t = p$ , then  $\overline{T(p, M)} - S(p, M) = \emptyset$  from  $T(p, M) \subset S(p, M)$ .

(d) implies (a): Let  $t = Bd S(p, M)$ . Then  $t \in \overline{T(p, M)}$  by Lemma 1.

If  $t \neq p$ , by assertion of (d), there exists a unique separation

$$M-t = \{M-\overline{S(p, M)}\} \cup S(p, M), \text{ where } M-\overline{S(p, M)} \ni p.$$

Hence,  $S(p, M)$  is a connected set. Therefore, we can assume  $t = p$ . Then, there exists a component  $K$  of  $S(p, M)$  such that

$$(4) \quad \overline{T(p, M)} - K \ni t.$$

Since  $T(p, M) \subset \overline{S(p, M)}$  and  $t \notin T(p, M)$ ,

$$(5) \quad T(p, M) - K = T(p, M) \cap \{S(p, M) - K\}.$$

Let us assume that  $S(p, M)$  is not connected. Then, there exists a component  $L$  of  $S(p, M)$  different from  $K$ . It follows from (5) that

$$(6) \quad \overline{T(p, M)} - K \supset L \cap \overline{T(p, M)}.$$

By (4) and (6), there exist neighbourhoods  $U, V$  of  $t$  and a connected set  $C$  satisfying  $U \supset C \supset V \ni t$  and

$$(7) \quad U \cap L \cap T(p, M) = \emptyset.$$

There exists a point  $z \in V \cap L$  by  $Bd L = t$ . Let  $H_2(q)$  be a connected sect of  $M$  not containing  $p$  with  $H_2(q) \ni z$ , where  $q \in T(p, M)$ . To see  $q \in L$ , let  $q \notin L$ . Then  $H_2(q) = L$ , because  $L$  is a component of  $S(p, M)$ . Hence  $t = Bd L = Bd H_2(q) = q$ . This contradiction implies  $q \in L$ . By (7) and  $q \in L \cap T(p, M)$ ,  $C \ni q$ . Since  $C \cap H_2(q) \ni z$ ,  $H_2(q) \supset C \ni t$ . This contradiction completes the proof.

**Theorem 4.** *Let  $M$  be a space. In order that all non-empty  $S(p, M)$  ( $p \in M$ ) be connected, it is necessary and sufficient that  $M$  have one and only one cut point by which  $M$  is uniquely separated.*

*Proof.* Sufficiency: Let  $t$  be the unique cut point of the space  $M$ , and  $M-t = M_1(t) \cup M_2(t)$  the separation by  $t$ . We have  $p \neq t$  for non-empty  $S(p, M)$  ( $p \in M$ ). Hence, if  $p$  is such a point belonging to  $M_1(t)$ , then  $S(p, M) = M_2(t)$  is a connected set.

Necessity: Suppose that  $M$  contains two distinct cut points  $q_1, q_2$ . Let  $M_1$  be a component of  $M-q_1$  containing  $q_2$ ,  $M_2 = M-(M_1 \cup q_1)$ ,  $N_1$  a component of  $M-q_2$  containing  $q_1$ , and  $N_2 = M-(N_1 \cup q_2)$ . Then there exist two separations such that

$$(8) \quad M-q_1 = M_1 \cup M_2, \quad M-q_2 = N_1 \cup N_2,$$

where  $q_2 \in M_1$ ,  $q_1 \in N_1$ . Then

$$M_1 \cap N_1 \neq \emptyset \text{ and } M_2 \cap N_2 = \emptyset.$$

To see that  $M_1 \cap N_1$  has a cut point of  $M$ , let us assume that  $M_1 \cap N_1$  has no cut point of  $M$ . Let  $y$  be a point of  $M_1 \cap N_1$  and let  $x$  be a point of  $S(p, M) - (M_2 \cup N_2)$ . Then there exist a cut point  $t$  of  $M$  and a separation

$$M - t = H_1(t) \cup H_2(t), \text{ where } y \in H_1(t), x \in H_2(t).$$

If  $q_1 = t$ , then

$$H_1(t) \cup H_2(t) = M_1 \cup M_2.$$

Hence  $x \in H_2(t) \subset M_2$  by the definition of  $M_2$ . This implies  $q_1 \neq t$ . Then, we have  $t \in M_1$  by  $q_1 \in H_1(t) \cup H_2(t)$ . Analogously,  $t \in N_1$ . Therefore, we obtain that  $S(y, M) = M_2 \cup N_2$  because  $M_1 \cap N_1$  has no cut point. This contradicts the connectedness of  $S(y, M)$ .

Now, for an arbitrary separation

$$M - u = M_1(u) \cup M_2(u),$$

where  $u$  is any cut point of  $M$  in  $M_1 \cap N_1$ , we have

$$(9) \quad \bar{M}_2 \subset M_1(u) \text{ or } \bar{M}_2 \subset M_2(u)$$

and

$$(10) \quad \bar{N}_2 \subset M_1(u) \text{ or } \bar{N}_2 \subset M_2(u).$$

On the other hand, by the connectedness of  $S(p, M)$  for each point  $p \in M$ , for a cut point  $u$  of  $M$  in  $M_1 \cap N_1$  there exists no separation

$$M - u = M_1(u) \cup M_2(u)$$

which satisfies

$$(11) \quad M_1(u) \supset M_2 \text{ and } M_2(u) \supset N_2$$

or

$$(12) \quad M_1(u) \supset N_2 \text{ and } M_2(u) \supset M_2.$$

In fact, if for any cut point  $u \in M_1 \cap N_1$  there exists a separation

$$M - u = M_1(u) \cup M_2(u)$$

satisfying (11) or (12), then  $M_1(u) \supset M_2$  and  $M_2(u) \supset N_2$ . This is contrary to the connectedness of  $S(u, M) \subset M - u$ .

Now, let  $z$  be a cut point of  $M$  in  $M_1 \cap N_1$  with the following separation

$$M-z = M_1(z) \cup M_2(z).$$

Here, from (9)~(12) we may assume that  $M_1(z) \supset \overline{M_2 \cup N_2}$ . Then

$$(13) \quad M_2(z) \subset S(q_1, M).$$

We shall see  $q_2 \notin S(q_1, M)$ . If not, there exist a cut point  $t$  and a separation

$$M-t = K_1(t) \cup K_2(t), \text{ where } K_1(t) \ni q_1, K_2(t) \ni q_2.$$

It follows that  $K_1(t) \supset \overline{M_2}$  and  $K_2(t) \supset \overline{N_2}$  by  $M_1 \cap N_1 \ni t$ . This contradicts (11) or (12). By (8) and the connectedness of  $S(q_1, M)$ , we have then

$$(14) \quad N_2 = S(q_1, M).$$

However, since  $N_2 \subset M_1(z)$ , (14) contradicts (13). This implies that  $M$  has a single cut point  $t$  of  $M$ . Hence, we have  $Bd S(p, M) = t$  for each  $p$  in  $M-t$  (Lemma 1) and  $M$  is uniquely separated by  $t$  (Theorem 3). This completes the proof.

#### REFERENCES

- [1] G.T. WHYBURN: Analytic Topology, Amer. Math. Soc. Coll. Publ. **28**, 1942.
- [2] R.L. MOORE: Foundation of Point Set Theory, Amer. Math. Soc. Coll. Publ. **13**, 1962.

DEPARTMENT OF MATHEMATICS,  
OKAYAMA UNIVERSITY

(Received June 17, 1971)