

ON s -UNITAL RINGS

Dedicated to Professor Mikao Moriya on his 70th birthday

HISAO TOMINAGA

The present paper attempts to generalize several results in [10], [21], [22] and [24] obtained for rings with identity. In fact, we can prove similar ones for left (and right) s -unital rings, where a ring $R (\neq 0)$ is called a left s -unital ring if $Ra \ni a$ for any $a \in R$. Needless to say, the class of left s -unital rings includes those of rings with identity and of regular rings. In [6], [18] and [23] we treated with left s -unital rings in the connection with regular rings. In the present paper, our attention will be directed towards the classes of left V -rings, fully left idempotent rings, and of almost commutative rings, those which are closely related to the class of regular rings. §1 contains a fundamental proposition, a characterization of prime ideals of a left s -unital ring in terms of its right modules as in Beachy [3], and a slight generalization of a result of Hansen [13]. The material of §2 comes from Fisher [10], Michler-Villamayor [21], Ramamurthi [22] and Yue Chi Ming [25], and left V -rings will be concerned in regular rings, left p - V -rings and fully left idempotent rings. In §§3 and 4, almost all results of Wong [24] will be carried over to s -unital rings.

For future reference, $R (\neq 0)$ will represent always a ring (with or without identity), and C the center of R . The Jacobson radical and the prime radical of R will be denoted by $J(R)$ and $P(R)$, respectively. As for other notations, we follow [18] and [23].

1. s -unital rings. A left R -module $M \neq 0$ is defined to be s -unital if $Ru \ni u$ for any $u \in M$. For instance, every irreducible left R -module is s -unital. Needless to say, if ${}_R M$ is s -unital then it is unital, and in case R contains 1 these notions are identical. We can define similarly an s -unital right R -module.

Theorem 1. *If $M (\neq 0)$ is a left R -module then the following are equivalent :*

- 1) ${}_R M$ is s -unital.
- 2) For any $u_1, \dots, u_n \in M$ there exists an element $e \in R$ such that $eu_i = u_i$ ($i=1, \dots, n$).

3) For any positive integer n , every $(R)_n$ -submodule of the direct sum ${}^{(n)}M$ of n copies of M is of the form ${}^{(n)}N$ with some ${}_R N \subseteq {}_R M$, where $(R)_n$ denotes the $n \times n$ matrix ring over R .

Proof. 1) \Leftrightarrow 2). Assume that ${}_R M$ is s -unital. Choose an element $e_n \in R$ such that $e_n u_n = u_n$, and set $v_i = u_i - e_n u_i$ ($i=1, \dots, n-1$). By induction method, there exists an element $e' \in R$ such that $e' v_i = v_i$ ($i=1, \dots, n-1$). Then, one will easily see that $e = e' + e_n - e' e_n$ is an element with the property requested in 2). The converse is trivial.

1) \Leftrightarrow 3). Given $a \in R$, $E_{ij}(a)$ will denote the element of $(R)_n$ with a in the (i, j) -position and zeros elsewhere. If $u_1, \dots, u_n \in M$ then

$$E_{11}(a) \begin{pmatrix} u_1 \\ \cdot \\ \cdot \\ \cdot \\ u_n \end{pmatrix} = \begin{pmatrix} au_1 \\ 0 \\ \cdot \\ \cdot \\ 0 \end{pmatrix},$$

whence we can easily see that 1) implies 3). The converse is also easy

by the fact that $\begin{pmatrix} Ru + Zu \\ Ru \\ \cdot \\ \cdot \\ Ru \end{pmatrix}$ is an $(R)_n$ -submodule of ${}^{(n)}M$ for any $u \in M$.

If ${}_R R$ (resp. R_R) is s -unital, R is said to be *left* (resp. *right*) s -unital. To be easily seen, every (non-zero) homomorphic image of a left s -unital ring is left s -unital, and any regular ring is left and right s -unital. (In Ramamurthi [22], a left s -unital ring is cited as a *left D-regular ring*.)

Corollary 1. *If R is left s -unital then so is $(R)_n$, and conversely.*

Proof. If $A = (a_{ij})$ is an arbitrary element of $(R)_n$, then by Theorem 1 there exists an element $e \in R$ such that $ea_{ij} = a_{ij}$ ($i, j=1, \dots, n$), whence it follows $\text{diag}\{e, \dots, e\} \cdot A = A$. Conversely, if $Ra \neq a$ then $(R)_n \cdot \text{diag}\{a, \dots, a\}$ does not contain $\text{diag}\{a, \dots, a\}$.

Proposition 1 (cf. [2, Proposition 5]). *Let τ be a non-zero right ideal of R . Then the following are equivalent:*

- 1) τ is a left s -unital ring.
- 2) $\tau \cap I = \tau I$ for any left ideal I of R .

If R is right s -unital then 1) is also equivalent to the following:

3) $\tau M \cap N = \tau N$ for any left R -modules ${}_R N \subseteq {}_R M$.

(In case R contains 1, it is known that 1) is nothing but to say that $(R/\tau)_R$ is flat (see for instance [19, Proposition 3, p. 133]).)

Proof. 1) \iff 2) is easy, and in case R is right s -unital 2) is obviously a special case of 3).

1) \implies 3). Let $u = a_1 u_1 + \dots + a_n u_n$ ($a_i \in \tau$, $u_i \in M$) be an arbitrary element of $\tau M \cap N$, and choose $e \in \tau$ with $ea_i = a_i$ for all i (Theorem 1). Then $u = ea_1 u_1 + \dots + ea_n u_n = eu \in \tau N$.

The next will play occasionally an important role in our subsequent study.

Proposition 2. *Let R be a left (resp. right) s -unital ring.*

(1) *If α is a proper ideal of R then α is contained in a proper prime ideal.*

(2) *Let R'/R be a ring extension. If α' is an ideal of R' and $\alpha' \cap R \neq R$ then there exists a maximal left (resp. right) ideal m' of R' such that $m' \supseteq \alpha'$ and $m' \cap R \neq R$. Especially, if α is a proper ideal of R then α is contained in a maximal left (resp. right) ideal of R (cf. [23, Lemma 1 (a)]).*

Proof. (1) Let $r \in R \setminus \alpha$, and choose $e \in R$ such that $r = er$. Then $E = \{e^i \mid i = 1, 2, \dots\}$ is an m -system excluding α . If $\mathfrak{p} \supseteq \alpha$ is an ideal of R which is maximal with respect to excluding E , then \mathfrak{p} is a proper prime ideal.

(2) Let $r \in R \setminus (\alpha' \cap R)$, and choose $e \in R$ such that $r = er$. By Zorn's lemma, there exists a maximal member m' in the family of left ideals b' of R' with $b' \supseteq \{x' \in R' \mid x'r \in \alpha'\} (\supseteq \alpha')$ and $b' \not\ni e$. Obviously $m' \cap R \neq R$, and one will easily see that m' is a maximal left ideal of R' .

For a right R -module M_R , we set $\tau(M_R) = \sum_r fM$ ($f \in \text{Hom}(M_R, R_R)$) and $\text{Ann}(M_R) = \{x \in R \mid Mx = 0\}$. To be easily seen, $\tau(M_R)$ is an ideal of R and $\text{Ann}(M_R) \subseteq \text{Ann}(\tau(M_R)_R)$.

Now, let M_R and M'_R be non-zero right R -modules. If for each $u \neq 0$ in M there exists $f \in \text{Hom}(M_R, M'_R)$ such that $fu \neq 0$, then we write $M_R > M'_R$. If $M_R > M'_R$ and $M'_R > M_R$, then we write $M_R \sim M'_R$. It is easy to see that the relations $>$ and \sim are transitive. Obviously, $M_R > R_R$ is nothing but to say that M_R is torsionless, and then we have $\text{Ann}(M_R) = \text{Ann}(\tau(M_R)_R)$. If M_R is faithful then $R_R > M_R$, and in case R is left s -unital the converse is also true.

In what follows, we shall present a characterization of proper prime ideal of a left s -unital ring in terms of its right modules. If R is a prime ring and $M_R > R_R$ then $\tau(M_R)$ is non-zero and $\text{Ann}(M_R) = \text{Ann}(\tau(M_R)_R) = 0$, namely, M_R is faithful. Conversely, if every torsionless right R -module is faithful then R is seen to be prime. Hence, for a left s -unital ring R , we see that R is prime if and only if $M_R > R_R$ implies always $M_R \sim R_R$.

Theorem 2 (cf. [3, Theorem 2]). *If \mathfrak{p} is a proper ideal of a left s -unital ring R then the following are equivalent :*

- 1) \mathfrak{p} is a prime ideal.
- 2) $M_R > (R/\mathfrak{p})_R$ implies always $M_R \sim (R/\mathfrak{p})_R$.

Proof. If $M_R > (R/\mathfrak{p})_R$ then $\text{Ann}(M_R) \supseteq \text{Ann}((R/\mathfrak{p})_R) = \mathfrak{p}$, and so M_R may be regarded as $M_{R/\mathfrak{p}}$. Hence, R/\mathfrak{p} is a prime ring if and only if $M_R \sim (R/\mathfrak{p})_R$ for any $M_R > (R/\mathfrak{p})_R$.

Corollary 2 (cf. [3, Theorem 3]). *Let R be a left s -unital ring. If $N_R (\neq 0)$ is a unital module then the following are equivalent :*

- 1) $M_R > N_R$ implies always $M_R \sim N_R$.
- 2) $N_R \sim (R/\mathfrak{p})_R$ for a proper prime ideal \mathfrak{p} .

Proof. 1) \Rightarrow 2). Let $\mathfrak{p} = \text{Ann}(N_R) (\neq R)$. Since $N_{R/\mathfrak{p}}$ is faithful, we have $(R/\mathfrak{p})_{R/\mathfrak{p}} > N_{R/\mathfrak{p}}$, and hence $(R/\mathfrak{p})_R \sim N_R$. If $M_R > (R/\mathfrak{p})_R$ then $M_R > N_R$, and $M_R \sim N_R \sim (R/\mathfrak{p})_R$, whence it follows that \mathfrak{p} is a prime ideal (Theorem 2).

2) \Rightarrow 1). Since $M_R > N_R \sim (R/\mathfrak{p})_R$ and \mathfrak{p} is prime, Theorem 2 shows that $M_R \sim (R/\mathfrak{p})_R \sim N_R$.

As was shown in [13], every left Noetherian, left s -unital ring has a left identity. The next is a slight generalization of the result.

Theorem 3. *If a left Goldie ring R is left s -unital then R contains a left identity.*

Proof. To be easily seen, the left singular ideal $Z_l(R)$ is contained in $P(R)$ that is nilpotent by Lanski's theorem (cf. [16, p. 24]). By [9, Theorem 1.3], $R/Z_l(R)$ satisfies the maximum condition for right annihilators. Then $R/Z_l(R)$ has a left identity by [14, Proposition 2.1], and hence the semi-prime ring $R/P(R)$ has the identity. Now, we shall proceed by the induction with respect to the nilpotency index n of $P(R)$. The case $n=1$ is obvious by the above. Assume $n > 1$. Since $R/P(R)^{n-1}$ has a left identity by the induction hypothesis and $R/P(R)$ has the identity,

a result of Herstein (cf. [15, p. 31]) shows that R has a left identity.

Corollary 3. *If R is left s -unital then the following are equivalent :*

- 1) R is a left Artinian ring.
- 2) R is a left Noetherian π -regular ring.
- 3) R is a fully left Goldie π -regular ring.

Proof. If R is left Artinian then R is left Noetherian by Hopkins' theorem (cf. [17, Theorem 34, p. 134]). Moreover, R being of bounded index, R is π -regular by [1, Theorem 5]. Since 2) implies 3) obviously, it remains only to prove that 3) implies 1). As was claimed in the proof of Theorem 3, $P(R)$ is nilpotent and $\bar{R} = R/P(R)$ has the identity. Now, let \bar{a} be an arbitrary regular element of \bar{R} , and $\bar{a}^n \bar{x} \bar{a}^n = \bar{a}^n$. Then, $\bar{a}^n (1 - \bar{x} \bar{a}^n) = 0$ implies $\bar{x} \bar{a}^n = 1$, and similarly $\bar{a}^n \bar{x} = 1$. Hence, every regular element of \bar{R} is a unit, which means that \bar{R} coincides with its left quotient ring that is Artinian semiprimitive. Recalling here that $R/P(R)^{k+1}$ is a left s -unital, left Goldie ring, one will easily see that ${}_{\bar{R}}(P(R)^k/P(R)^{k+1})$ is completely reducible and of finite length. It follows therefore that ${}_R R$ has a composition series.

Corollary 4. *Let R be a left s -unital, fully left Goldie ring whose prime factor rings are π -regular. If α is an ideal of R and ${}_{R\alpha} R$ is of finite length, then ${}_R \alpha$ is of finite length.*

Proof. To our end, it suffices to prove the assertion for a minimal ideal α . Obviously, $l(\alpha)$ is a prime ideal of R and $S = R/l(\alpha)$ is Artinian simple by Corollary 3. Since R is left Goldie and ${}_S \alpha$ is completely reducible, ${}_R \alpha$ is of finite length.

The next is perhaps in the same vein as Corollary 4, and can be proved in the same way as in the proof of [20, Proposition].

Corollary 5. *Let R be a left s -unital, left Noetherian ring. If α is an ideal of R and ${}_{\alpha} R$ is of finite length, then ${}_R \alpha$ is of finite length, too.*

Remarks. (1) Every s -unital left R -module is a homomorphic image of a direct sum of copies of ${}_R R$.

(2) Let M be an s -unital left R -module over a left s -unital ring R . We consider the map $f: M \longrightarrow R \otimes_R M$ defined by $u \longmapsto e' \otimes u$, where $e'u = u$. If $e''u = u$ ($e'' \in R$) then there exists an element $e \in R$ such that $ee' = e'$ and $ee'' = e''$ (Theorem 1) and we have $e' \otimes u = ee' \otimes u = e \otimes e'u$

$= e \otimes e''u = e'' \otimes u$. Hence, f is well-defined and is an R -homomorphism. Now, let $\sum_i a_i \otimes u_i$ be an arbitrary element of $R \otimes_R M$. Again by Theorem 1, we can find an element $a \in R$ such that $aa_i = a_i$ for all i . We have then $(\sum_i a_i u_i) f = a \otimes \sum_i a_i u_i = \sum_i aa_i \otimes u_i = \sum_i a_i \otimes u_i$. This proves that ${}_R R \otimes_R M$ is canonically isomorphic to ${}_R M$. Similarly, if R is commutative then we can prove the same for any s -unital module ${}_R M$.

(3) An s -unital module M_R will be defined to be s -flat if for each pair of s -unital left R -modules $A \subseteq B$ (with the inclusion map $\iota: 1 \otimes \iota: M \otimes_R A \rightarrow M \otimes_R B$ is a monomorphism. As a consequence of (2), one will easily see that if R is left and right s -unital then R_R is s -flat. Moreover, we can prove the following: Let R be a left and right s -unital ring, and \mathfrak{l} a left ideal of R . If M_R is s -flat then $M \otimes_R \mathfrak{l}$ is canonically isomorphic to $M\mathfrak{l}$.

2. V-rings. An s -unital left R -module M is defined to be s -injective if M has the property that for each pair of s -unital left R -modules $A \subseteq B$ each $f \in \text{Hom}({}_R A, {}_R M)$ can be extended to an element of $\text{Hom}({}_R B, {}_R M)$. If ${}_R M$ is s -injective then ${}_R M < \bigoplus {}_R M'$ for any s -unital ${}_R M' \supseteq {}_R M$. Moreover, the proof of [8, Theorem 1.6] enables us to obtain the following:

Proposition 3 (Baer Criterion). *Let R be a left s -unital ring, and M an s -unital left R -module. Then ${}_R M$ is s -injective if and only if for each left ideal \mathfrak{l} of R each $f \in \text{Hom}({}_R \mathfrak{l}, {}_R M)$ can be extended to an element of $\text{Hom}({}_R R, {}_R M)$.*

An s -unital left (resp. right) R -module M is called a V -module if every R -submodule of M is an intersection of maximal R -submodules. If ${}_R R$ (resp. R_R) is a V -module, R is called a *left* (resp. *right*) V -ring (cf. [5]). As was mentioned in [18, Remark], we obtain the following which corresponds to [21, Theorem 2.1]:

Theorem 4. *The following are equivalent:*

- 1) R is a left V -ring.
- 2) R is left s -unital and every irreducible left R -module is s -injective.
- 3) R is left s -unital and every s -unital left R -module is a V -module.
- 4) R is left s -unital, and for any s -unital left R -module M the intersection of all maximal R -submodules is 0; $\text{rad } {}_R M = 0$.
- 5) For any positive integer n , $(R)_n$ is a left V -ring.

Proof. First, we shall prove the equivalence of 1)–4). Obviously, 4) \Leftrightarrow 3) \Rightarrow 1).

2) \Rightarrow 4). Let M be an arbitrary s -unital left R -module. If $0 \neq u \in$

M , then there exists an R -submodule Y of M which is maximal with respect to $Y \not\supseteq u$. Let S be the set of R -submodules of M properly containing Y , and $D = \bigcap_{X \in S} X (\ni u)$. Since D/Y is an irreducible R -module, by 2) there exists an R -submodule K of M containing Y such that $M/Y = D/Y \oplus K/Y$. Then $u \notin K$, and hence $Y = K$, namely, $M = D$. This means that Y is a maximal R -submodule of M and $\text{rad } {}_R M = 0$.

1) \implies 2). Let M be an irreducible left R -module, and I a left ideal of R . If f is a non-zero element of $\text{Hom}({}_R I, {}_R M)$, then $I' = \text{Ker } f \subset I$. By 1), there exists a maximal left ideal m such that $m \supseteq I'$ and $m \not\supseteq I$. Since ${}_R M \cong {}_R(I/I')$ is irreducible and $I \supset m \cap I \supseteq I'$, we have $m \cap I = I'$. Now, taking this into mind, we can well-define an extension $g \in \text{Hom}({}_R R, {}_R M)$ of f by $l + m \mapsto lf (l \in I, m \in m)$. Hence ${}_R M$ is s -injective by Proposition 3.

Next, we shall prove 1) \implies 5) \implies 4).

1) \implies 5). The direct sum $R^{(n)}$ of n copies of R is an s -unital left R -module (Theorem 1), and we have seen that ${}_R R^{(n)}$ is a V -module. Again by Theorem 1, every $(R)_n$ -submodule of $(R)_n = {}^{(n)}(R^{(n)})$ is of the form ${}^{(n)}N$ with some ${}_R N \subseteq {}_R R^{(n)}$. Since ${}_R R^{(n)}$ is a V -module, $N = \bigcap_{\alpha} M_{\alpha}$ with maximal submodules ${}_R M_{\alpha} \subseteq {}_R R^{(n)}$. Hence ${}^{(n)}N = \bigcap_{\alpha} {}^{(n)}M_{\alpha}$, and $(R)_n$ is a left V -ring.

5) \implies 4). Again by Theorem 1, given an s -unital ${}_R M$, the left $(R)_n$ -module ${}^{(n)}M$ is s -unital and $\text{rad } {}_{(R)_n} {}^{(n)}M = 0$, whence it follows $\text{rad } {}_R M = 0$.

A left R -module M is said to be p -injective if for any principal left ideal $(a|$ of R and $f \in \text{Hom}({}_R(a|, {}_R M)$ there exists an element $u \in M$ such that $xf = xu$ for all $x \in (a|$. As was noted in [6], R is regular if and only if every left R -module is p -injective (cf. also [25]). In connection with Theorem 3, a left s -unital ring R is defined to be a *left p - V -ring* if every irreducible left R -module is p -injective. We can define a *right p - V -ring* in an obvious way. In case R contains 1, a left V -ring is a left p - V -ring. More generally we have

Proposition 4. *If R is a right s -unital, left V -ring then it is a left p - V -ring.*

Proof. Let ${}_R M$ be irreducible, and $(a| (= Ra)$ an arbitrary principal left ideal of R . Choose an element $e \in R$ with $ae = a$. If $f \in \text{Hom}({}_R(a|, {}_R M)$ and $g \in \text{Hom}({}_R R, {}_R M)$ is an extension of f , then for any $x \in R$ there holds $(xa)f = (xa)g = (xae)g = xa \cdot eg$.

If every left (resp. right) ideal of R is idempotent, R is said to be *fully left* (resp. *right*) *idempotent*. (In [22], a fully left idempotent ring is cited as a *left weakly regular ring*.) On the other hand, R is said to be *fully idempotent* if every ideal of R is idempotent.

Proposition 5. (1) *The following are equivalent :*

- 1) *R is fully left idempotent.*
- 2) $(Ra)^2 \ni a$ for any $a \in R$.
- 3) *For each pair of left ideals $\mathfrak{l} \subseteq \mathfrak{l}'$ of R, there holds $\mathfrak{l}'\mathfrak{l} = \mathfrak{l}$.*
- 4) *For any positive integer n, $(R)_n$ is fully left idempotent.*

If R is right s-unital then 1) is also equivalent to each of the following :

- 5) *For each ideal \mathfrak{a} and each left ideal \mathfrak{l} of R there holds $\mathfrak{a} \cap \mathfrak{l} = \mathfrak{a}\mathfrak{l}$.*
- 6) *For each ideal \mathfrak{a} of R and each pair of left R-modules ${}_R N \subseteq {}_R M$ there holds $\mathfrak{a}M \cap N = \mathfrak{a}N$.*

(2) *The following are equivalent :*

- 1) *R is fully idempotent.*
- 2) $(RaR)^2 \ni a$ for any $a \in R$.
- 3) *Every ideal of R is semiprime.*
- 4) *For each pair of ideals $\mathfrak{a}, \mathfrak{a}'$ of R there holds $\mathfrak{a} \cap \mathfrak{a}' = \mathfrak{a}\mathfrak{a}'$.*
- 5) *For any positive integer n, $(R)_n$ is fully idempotent.*

Proof. The assertion (2) is given in [7]. Concerning (1), the equivalence of 1)–3) is given in [22, Proposition 1] and 4) \Rightarrow 1) is trivial. Moreover, the latter part will be obvious by Proposition 1.

1) \Rightarrow 4). We shall modify slightly the proof of [12, Theorem 4].

At first, we consider the case $n = 2$. Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ be an arbitrary element of $\mathfrak{R} = (R)_2$. If $a = \sum_i w_i a w'_i a$ ($w_i, w'_i \in R$), then $A - X = \begin{pmatrix} 0 & b' \\ c & d \end{pmatrix}$ for $X = \sum_i \begin{pmatrix} w_i & 0 \\ 0 & 0 \end{pmatrix} A \begin{pmatrix} w'_i & 0 \\ 0 & 0 \end{pmatrix} A$. Next, if $d = \sum_j x_j d x'_j d$ ($x_j, x'_j \in R$), then $A - X - Y = \begin{pmatrix} 0 & b' \\ c' & 0 \end{pmatrix}$ for $Y = \sum_j \begin{pmatrix} 0 & 0 \\ 0 & x_j \end{pmatrix} (A - X) \begin{pmatrix} 0 & 0 \\ 0 & x'_j \end{pmatrix} (A - X)$. Finally, if $b' = \sum_k y_k b' y'_k b'$ and $c' = \sum_k z_k c' z'_k c'$ ($y_k, y'_k, z_k, z'_k \in R$) then $A - X - Y = \sum_k \begin{pmatrix} y_k & 0 \\ 0 & z_k \end{pmatrix} (A - X - Y) \begin{pmatrix} 0 & z'_k \\ y'_k & 0 \end{pmatrix} (A - X - Y)$. We obtain therefore $A = (A - X - Y) + Y + X \in (\mathfrak{R}(A - X - Y))^2 + (\mathfrak{R}(A - X))^2 + (\mathfrak{R}A)^2 = (\mathfrak{R}A)^2$, namely, \mathfrak{R} is fully left idempotent.

Since $(R)_{2^k} \cong (R_{2^{k-1}})_2$, one will easily see that $(R)_{2^k}$ is fully left idempotent. Given arbitrary n , we choose k so that $2^k \geq n$. If $A \in (R)_n$, we choose $A' \in (R)_{2^k}$ with A in the upper left-hand corner and zeros elsewhere. Now, $A' \in ((R)_{2^k} A')^2$ and a brief computation gives $A \in ((R)_n A)^2$.

Proposition 6. *Every left p-V-ring is fully left idempotent, and so every right s-unital, left V-ring is fully left idempotent.*

Proof. If not, there exists a non-zero element $a \in R$ such that $(a|^2 \neq (a| (=Ra)$. Let m be a maximal member in the family of left ideals l of R such that $(a|^2 \subseteq l \subset (a|$. Since the irreducible left R -module $(a|/m$ is p -injective, there exists an element $b \in (a|$ such that $x + m = xb + m$ for all $x \in (a|$. But this implies a contradiction $(a| = m$. The latter part is evident by Proposition 4.

As a direct consequence of Proposition 6 and [11, Theorem 1.1], we obtain the following :

Corollary 6 (cf. [10, Theorem 13]). *If R is a left V -ring then the following are equivalent :*

- 1) R is a regular ring.
- 2) R is right s -unital and every prime factor ring of R is a regular ring.

Proposition 7 (cf. [21], [22]). *Let R be fully left idempotent.*

- (1) R is right non-singular ; $Z_r(R) = 0$.
- (2) R is semiprimitive ; $J(R) = 0$.
- (3) If $a \in R$ is left regular then $R = RaR$.
- (4) C is a regular ring.¹⁾

Proof. (1) Let $z \in Z_r(R)$, and choose an element $y \in Z_r(R)$ such that $z = yz$ (Proposition 5 (1)). If $nz + zx$ (n an integer and $x \in R$) is an arbitrary element of $|z) \cap r(y)$, then $0 = y(nz + zx) = nyz + yzx = nz + zx$. Hence, $|z) \cap r(y) = 0$, which means $z = 0$.

(2) Let $z \in J(R)$, and choose $y \in J(R)$ such that $z = yz$. Since $\{xy - x | x \in R\} = R$, it follows $Rz = 0$, namely, $z = 0$.

(3) This is evident by $Ra = RaRa$.

(4) If c is an arbitrary element of C then $c \in (Rc)^2 = Rc^2$ by Proposition 5 (1). Hence, C is regular by [1, Lemma 1].

Theorem 5 (cf. [10, Theorem 14]). *If R is right s -unital then the following are equivalent :*

- 1) R is a left V -ring.

¹⁾ If R is fully idempotent then it is almost evident that C is still regular and the centroid \mathfrak{C} of R is commutative. Moreover, as was shown by R. Courter [Prcc. Amer. Math. Soc. 43 (1974), 293-295], \mathfrak{C} is a regular ring. In fact, given an arbitrary element γ of \mathfrak{C} , one will easily see that $R\gamma^2 = (R^2)\gamma^2 = (R\gamma)^2 = R\gamma$ and $R\gamma \cap \text{Ker } \gamma = (R\gamma \cap \text{Ker } \gamma)^2 = 0$. Hence, $R = R\gamma \oplus \text{Ker } \gamma$ and γ induces an automorphism of ${}_R R \gamma_R$. We can find then an element γ' of \mathfrak{C} such that $\gamma = \gamma^2 \gamma'$.

2) R is fully left idempotent and every left primitive factor ring of R is a left V -ring.

Proof. 1) \Rightarrow 2). This is a consequence of Proposition 6.

2) \Rightarrow 1). Let M be an irreducible left R -module, and I a left ideal of R . Let f be a non-zero element of $\text{Hom}({}_R I, {}_R M)$. Obviously $\alpha = \text{Ann}({}_R M)$ is a left primitive ideal of R . Noting that $\alpha \cap I = \alpha I$ (Proposition 5 (1)), one will easily see the map defined by $I + \alpha \ni l + a \mapsto lf$ ($l \in I, a \in \alpha$) is an extension of f in $\text{Hom}({}_R(I + \alpha), {}_R M)$. Now, the rest of the proof proceeds in the same way as for 1) \Rightarrow 2) of Theorem 4.

Corollary 7 (cf. [10, Corollary 15]). *If R is a regular ring then the following are equivalent :*

- 1) R is a left V -ring.
- 2) Every left primitive factor ring of R is a left V -ring.

A left (resp. right) s -unital ring is said to be *left* (resp. *right*) *semiartinian* if every s -unital left (resp. right) R -module contains an irreducible R -submodule.

Theorem 6 (cf. [10, Theorem 17]). *If R is left semiartinian then the following are equivalent :*

- 1) R is a regular ring.
- 2) R is fully idempotent.
- 3) R is fully left idempotent.
- 3') R is fully right idempotent.
- 4) R is a left p - V -ring.
- 4') R is a right p - V -ring.

When this is the case, R is right semiartinian.

Proof. 1) \Rightarrow 4) (resp. 4') \Rightarrow 3) (resp. 3') \Rightarrow 2). These are obvious by the remark mentioned before Proposition 4 and Proposition 6.

2) \Rightarrow 1). Let $S (\neq 0)$ be the left socle of R . If I is a left ideal of S then it is easy to see that $R I = R I \cdot R I \subseteq S I \subseteq I$, namely, I is a left ideal of R . (Note that R is semiprime.) Hence ${}_s S$ is completely reducible. Since S is also semiprime, each homogeneous component of ${}_s S$ is (non-trivial) simple and regular. Hence S is regular. Now, let $\mathfrak{m} (\supseteq S)$ be the maximal regular ideal of R (cf. [4]). Suppose $\mathfrak{m} \neq R$. Then R/\mathfrak{m} is fully idempotent and has non-zero left socle. By the above argument, we see that the maximal regular ideal of R/\mathfrak{m} is non-zero, which contradicts the maximality of \mathfrak{m} . We have seen thus $R = \mathfrak{m}$. Finally, noting that S coincides with the right socle of R , one will easily see that R has a right

socle sequence, namely, R is right semiartinian.

3. AC-rings. R is called an *AC-ring* (almost commutative ring) if for any proper prime ideal \mathfrak{p} of R and $a \notin \mathfrak{p}$ there exists x such that $ax \in C \setminus \mathfrak{p}$. Any P_1 -ring is obviously an AC-ring (cf. [6]), and the next will be easily seen (cf. [24, Theorem 1]).

Proposition 8. *Let R be an AC-ring.*

- (1) *Every homomorphic image of R is an AC-ring.*
- (2) *Every prime ideal of R is completely prime. In particular, R is a prime ring if and only if it is a domain.*
- (3) *Every semiprime ideal of R is completely semiprime. In particular, R is a semiprime ring if and only if it is a reduced ring.*
- (4) *For any proper prime ideal \mathfrak{p} of R and $a \notin \mathfrak{p}$ there exists y such that $ya \in C \setminus \mathfrak{p}$. (The notion of AC is right-left symmetric.)*

By Proposition 8 (3), the prime radical of an AC-ring coincides with the set of all nilpotent elements. If R is an AC-ring and $R \neq P(R)$ then $(R)_n$ cannot be an AC-ring for $n > 1$.

Proposition 9. *The following are equivalent :*

- 1) *R is a division ring.*
- 2) *$aR = R$ for any $a \neq 0$ in R .*
- 2') *$Ra = R$ for any $a \neq 0$ in R .*
- 3) *R is a (non-trivial) simple AC-ring.*
- 4) *R is a prime AC-ring with minimum condition on ideals.*
- 5) *R is a fully idempotent, prime AC-ring.*
- 6) *R is a regular, prime AC-ring.*

Proof. Obviously, 1) implies each of 2)—6) and 6) implies 5). Next, assume 2). Since R is strongly regular, there exists x such that $axa = a$ and $ax = xa$. By $axR = aR = R$, we see that the central idempotent ax is the identity of R , and 1) is obvious. Similarly, 1) \implies 2') \implies 1). Finally, assume one of 3)—5). For any $a \neq 0$ there exists x such that ax is a non-zero central element. Since R is a domain (Proposition 8), one will easily see $R = axR = aR$.

Corollary 9 (cf. [24, Theorem 3]). *The following are equivalent :*

- 1) *R is a finite direct sum of division rings.*
- 2) *R is a semiprime AC-ring with minimum condition on ideals.*
- 3) *R is a semiprimitive AC-ring with minimum condition on ideals.*

Proof. It suffices to prove $2) \Rightarrow 1)$. For any proper prime ideal \mathfrak{p} , R/\mathfrak{p} is a division ring (Proposition 9). Hence, R is a subdirect sum of division rings. As is well known, by the minimum condition on ideals, R is then a finite direct sum of division rings.

Theorem 7 (cf. [24, Theorem 2]). *If R is a left (resp. right) s -unital AC-ring and \mathfrak{n} is a submodule of R , then the following are equivalent :*

- 1) \mathfrak{n} is a maximal right (resp. left) ideal.
- 2) \mathfrak{n} is a maximal ideal.
- 3) \mathfrak{n} is a right (resp. left) primitive ideal.

Proof. Since R is left s -unital, $R^2 = R$ and any maximal ideal of R is a prime ideal. Moreover, if \mathfrak{n} is a right ideal of R then $(\mathfrak{n} : R) = \{x \in R \mid Rx \subseteq \mathfrak{n}\}$ coincides with the largest ideal contained in \mathfrak{n} .

$1) \Rightarrow 2)$. If $\mathfrak{a} = (\mathfrak{n} : R)$ ($\subseteq \mathfrak{n} \neq R$) is not maximal, then \mathfrak{a} is properly contained in a proper prime ideal \mathfrak{p} (Proposition 2 (1)). Evidently, there exists an element $a \in \mathfrak{n} \setminus \mathfrak{p}$, and $ax \in (\mathfrak{C} \cap \mathfrak{n}) \setminus \mathfrak{p}$ for some x . But this is impossible by $ax \in \mathfrak{a} \subseteq \mathfrak{p}$. This proves that \mathfrak{a} is a maximal ideal. Hence, R/\mathfrak{a} is a division ring (Proposition 9), and $\mathfrak{n} = \mathfrak{a}$.

$2) \Rightarrow 3)$. Since R/\mathfrak{n} is a division ring (Proposition 9), \mathfrak{n} is primitive.

$3) \Rightarrow 1)$. There exists a maximal right ideal \mathfrak{m} of R such that $\mathfrak{m} \supseteq \mathfrak{n}$ and $(\mathfrak{m}/\mathfrak{n} : R/\mathfrak{n}) = 0$. Since R/\mathfrak{n} is a left s -unital AC-ring, as was shown in $1) \Rightarrow 2)$, we obtain $\mathfrak{m}/\mathfrak{n} = (\mathfrak{m}/\mathfrak{n} : R/\mathfrak{n}) = 0$, i. e., $\mathfrak{m} = \mathfrak{n}$.

By Theorem 7, if R is a left (resp. right) s -unital AC-ring then $J(R)$ is the intersection of maximal ideals, and so a left (resp. right) s -unital semiprimitive AC-ring is a subdirect sum of division rings.

Theorem 8. *The following are equivalent :*

- 1) R is a strongly regular ring.
- 2) R is a regular AC-ring.
- 3) R is an AC-ring and a left (or right) p -V-ring.
- 4) R is a fully idempotent AC-ring.
- 5) R is an AC-ring whose ideals are semiprime.
- 6) R is a reduced ring such that R/\mathfrak{p} is regular (in fact a division ring) for any proper prime ideal \mathfrak{p} .
- 7) R is a reduced ring whose proper completely prime ideals are maximal left ideal.

Proof. $1) \Leftrightarrow 6) \Leftrightarrow 7)$ are given in [5] (and also in [11]), $1) \Rightarrow 2) \Rightarrow 3)$ are trivial, $3) \Rightarrow 4)$ by Proposition 6, and $4) \Rightarrow 6)$ is a consequence of Propositions 8 and 9. Finally, $4) \Leftrightarrow 5)$ is contained in Propo-

sition 5 (2).

Following [24], R is *primary* if every zero-divisor is nilpotent, and is *local* if it has exactly one maximal ideal.

Theorem 9 (cf. [24, Theorem 5]). (1) *If R is an AC-ring then the following are equivalent :*

- 1) *R is primary.*
- 2) *Every right zero-divisor is nilpotent.*
- 3) *Every left zero-divisor is nilpotent.*
- 4) *There exists a minimal prime ideal \mathfrak{p} of R which contains all zero-divisors.*

(2) *If R is a left s -unital AC-ring then the following are equivalent :*

- 1) *R has a unique prime ideal $\mathfrak{p} \neq R$.*
- 2) *R is local and $P(R) = J(R)$.*
- 3) *$R/P(R)$ is a division ring.*

Proof. (1) 2) \implies 3). Let $xy=0$, $y \neq 0$. If $x \notin P(R)$ then $x \notin \mathfrak{p}_0$ for some prime ideal \mathfrak{p}_0 . Choose $u \in R$ such that $ux \in C \setminus \mathfrak{p}_0$ (cf. Proposition 8). But, by 2), $0 = uxy = yux$ yields a contradiction $ux \in P(R)$. Similarly, we have 3) \implies 2). Obviously, $P(R)$ is a prime ideal.

1) \implies 2) \implies 4). Trivial.

4) \implies 1). It suffices to show that if x is non-nilpotent then $x \notin \mathfrak{p}$. To be easily seen, $T = \{x^k s \mid k \geq 0, s \in R \setminus \mathfrak{p}\} \cup \{x^k \mid k > 0\}$ is an m -system such that $x \in T$ and $0 \notin T$. Then there exists a prime ideal \mathfrak{p}_0 such that $\mathfrak{p}_0 \cap T = \emptyset$. Since \mathfrak{p} is a minimal prime ideal and $\mathfrak{p}_0 \subseteq R \setminus T \subseteq \mathfrak{p}$, we have $\mathfrak{p} = \mathfrak{p}_0 \not\ni x$.

(2) 1) \implies 2). Every maximal ideal of R is a prime ideal. If \mathfrak{p} is not maximal then it is properly contained in a proper prime ideal (Proposition 2 (1)), a contradiction.

2) \implies 3). Since $P(R) = J(R)$ is a unique maximal ideal, $R/P(R)$ is a division ring (Proposition 9).

3) \implies 1). Trivial.

4. Integral extensions of s -AC-rings. In [24], R with 1 is called an *SAC-ring* if for any proper ideal \mathfrak{a} and $x \notin \mathfrak{a}$ there exists y such that $xy \in C \setminus \mathfrak{a}$. However, in our present study, we shall employ a somewhat weaker (but right-left symmetric) definition: An AC-ring is called an *s -AC-ring* if for any non-prime ideal \mathfrak{a} and $x \notin \mathfrak{a}$ there holds $RxR \cap C \not\subseteq \mathfrak{a}$. To be easily seen, every s -AC-ring has the following property :

(*) For any proper ideal \mathfrak{a} and $x \notin \mathfrak{a}$ there holds $RxR \cap C \not\subseteq \mathfrak{a}$.

Any strongly regular ring is s -AC, and conversely any P_1 -ring with the property (*) is strongly regular.

For a while, we assume that R is a ring with the property (*). By a routine manner, we can show that an ideal α is prime (resp. semiprime) if and only if $\alpha \cap C$ is prime (resp. semiprime) in C . Accordingly, a ring is strongly regular if and only if it is an s -AC-ring whose center is regular (Theorem 8). We assume further that R'/R is a ring extension such that C is contained in the center C' of R' . Then we can easily see that if α' is a prime (resp. semiprime) ideal of R' then $\alpha' \cap R$ is a prime (resp. semiprime) ideal of R .

In what follows, R'/R will mean a ring extension, and C' the center of R' . R'/R is called a *left integral extension* if $C \subseteq C'$ and for each $x \in R'$ there exist a_0, \dots, a_{n-1} in R such that $x^n + a_{n-1}x^{n-1} + \dots + a_0 = 0$.

Concerning "going up" we have the following:

Theorem 10 (cf. [24, Theorem 7 and Corollary 2]). *Let R be an s -AC-ring, R' a left (or right) s -unital ring, and let R'/R be a left integral extension. If α' is an ideal of R' and \mathfrak{p} is a proper prime ideal of R containing $\alpha' \cap R$, then there exists a proper prime ideal \mathfrak{p}' of R' such that $\mathfrak{p}' \supseteq \alpha'$ and $\mathfrak{p}' \cap R = \mathfrak{p}$.*

Proof. Let M be the non-empty m -system $R \setminus \mathfrak{p}$, and $\mathfrak{p}' \supseteq \alpha'$ an ideal of R' which is maximal with respect to excluding M . Then \mathfrak{p}' is a proper prime ideal and $\mathfrak{p}' \cap R \subseteq \mathfrak{p}$. If $\mathfrak{p}' \cap R \subset \mathfrak{p}$ then there exists $c \in (C \cap \mathfrak{p}) \setminus (\mathfrak{p}' \cap R)$. Since $(cR' + \mathfrak{p}') \cap M \neq \emptyset$, $cx + \mathfrak{p}' = m$ with some $x \in R'$, $\mathfrak{p}' \in \mathfrak{p}'$, $m \in M$. Suppose $x^n + a_{n-1}x^{n-1} + \dots + a_0 = 0$ ($a_i \in R$). Then $0 = x^n c^n + a_{n-1}x^{n-1}c^n + \dots + a_0c^n = (m - \mathfrak{p}')^n + a_{n-1}(m - \mathfrak{p}')^{n-1}c + \dots + a_0c^n$. There exist therefore $r \in R$ and $q' \in \mathfrak{p}'$ such that $m^n + rc + q' = 0$. This shows $q' \in \mathfrak{p}' \cap R \subseteq \mathfrak{p}$, and hence $m^n \in \mathfrak{p}$, whence it follows a contradiction $m \in \mathfrak{p}$ (Proposition 8).

Corollary 9. *Let R be a left s -unital s -AC-ring, R' a left s -unital ring, and let R'/R be a left integral extension. If α is a proper ideal of R then $\alpha R'$ is a proper ideal of R' .*

Proof. By Proposition 2 (1), α is contained in a proper prime ideal \mathfrak{p} of R , and then there exists a proper prime ideal \mathfrak{p}' of R' such that $\mathfrak{p}' \cap R = \mathfrak{p}$ (Theorem 10). Hence $\alpha R' \subseteq \mathfrak{p}' \neq R'$. Next, to be easily seen, $R(\alpha \cap C) = \alpha$. It follows therefore $R'(\alpha R') = R'R(\alpha \cap C)R' \subseteq \alpha R'$.

Lemma 1. *Let R'/R be a left integral extension. If a completely prime ideal \mathfrak{p}' of R' is contained in a left ideal \mathfrak{v}' and $\mathfrak{v}' \cap R = \mathfrak{p}' \cap R$, then*

$n' = p'$.

Proof. Suppose there exists $x \in n' \setminus p'$. Let n be the smallest integer such that $x^n + a_{n-1}x^{n-1} + \dots + a_0 = p' \in p'$ ($a_i \in R$). This implies $a_0 \in n' \cap R = p' \cap R$ and $n > 1$. Since p' is completely prime, $(x^{n-1} + a_{n-1}x^{n-2} + \dots + a_1)x = p' - a_0 \in p'$ yields a contradiction $x^{n-1} + \dots + a_1 \in p'$.

Theorem 11 (cf. [24, Corollary 4]). *Let R be a right and left s -unital s -AC-ring, R' a left s -unital ring, and let R'/R be a left integral extension. Let p' be a completely prime ideal of R' . Then p' is a maximal left ideal if and only if $p' \cap R$ is a maximal ideal of R .*

Proof. Suppose $p' \cap R$ is a maximal ideal. By Proposition 2 (2), there exists a maximal left ideal m' of R' such that $m' \supseteq p'$ and $m' \cap R \neq R$. Since $p' \cap R$ is a maximal left ideal by Theorem 7, we have $m' \cap R = p' \cap R$, whence it follows $m' = p'$ (Lemma 1). Conversely, suppose p' is a maximal left ideal. We claim here $p' \cap R \neq R$. In fact, if $x \in R' \setminus p'$ and $x^n + a_{n-1}x^{n-1} + \dots + a_0 = 0$ ($a_i \in R$) then $p' \supseteq R$ gives a contradiction $x^n \in p'$. Now, suppose $p' \cap R$ is not maximal. Then $p' \cap R$ is properly contained in a proper prime ideal p_0 of R (Proposition 2 (1)), and we can find a proper prime ideal $p'_0 \supseteq p'$ of R' such that $p'_0 \cap R = p_0$ (Theorem 10). Since p'_0 has to be equal to p' , we have a contradiction $p' \cap R = p'_0 \cap R = p_0$.

Theorem 12 (cf. [24, Theorem 9]). *Let R be a left and right s -unital s -AC-ring, R' a left and right s -unital reduced ring, and let R'/R be a left integral extension. Then, R is regular if and only if so is R' .*

Proof. If R is (strongly) regular then every proper prime ideal of R is a maximal left ideal (Theorem 8). By the proof of Theorem 11, for any proper completely prime ideal p' of R' , $p' \cap R$ is a proper prime ideal of R , and so it is a maximal left ideal. Hence p' is a maximal left ideal by Theorem 11, and again by Theorem 8 R' is a regular ring. Conversely, if R' is a regular ring, then for any proper prime ideal p of R there exists a proper prime ideal p' of R' such that $p' \cap R = p$ (Theorem 10) and p' is a maximal left ideal (Theorem 8). Hence, p is a maximal ideal by Theorem 11, and so it is a maximal left ideal (Theorem 7). Theorem 8 proves therefore that R is regular.

Theorem 13 (cf. [24, Theorem 10]). *Let R'/R be a left integral extension. If R is strongly regular then for each $x \in R'$ there exist $y \in R'$ which can be expressed as a (left) polynomial in x over $R^1 = R + Z$ and a natural number n such that $yx^{n+1} = x^n$.*

Proof. Let $A(x) = \{p(x) \mid p(x) \text{ is a monic polynomial of positive degree in } x \text{ over } R \text{ such that } p(x)x^m = 0 \text{ for some } m\}$. In $A(x)$ we choose $p(x) = x^k + a_{k-1}x^{k-1} + \cdots + a_1x + a_0$ of the least degree; $p(x)x^{n-1} = 0$ ($n > 1$). By [1, Lemma 1], there exists (uniquely) an element $a \in R$ such that $a_0a = aa_0$, $a_0^2a = a_0$ and $a^2a_0 = a$. Obviously, $e = a_0a$ is a central idempotent with $ea_0 = a_0$ and $ea = a$. If $k = 1$ then $x^n + a_0x^{n-1} = 0$ implies $0 = x^n + a_0x^{n-1} - e(x^n + a_0x^{n-1}) = x^n - ex^n$, i. e., $ex^n = x^n$. Hence, $0 = a(x^n + a_0x^{n-1})x = ax^{n+1} + ex^n = ax^{n+1} + x^n$, whence it follows $-ax^{n+1} = x^n$. Next, we shall consider the case $k > 1$, and set $x_0 = x - ex$. Since $0 = p(x)x^{n-1} - ep(x)x^{n-1} = (x^k + a_{k-1}x^{k-1} + \cdots + a_1x)x^{n-1} - e(x^k + a_{k-1}x^{k-1} + \cdots + a_1x)x^{n-1} = (x_0^k + a_{k-1}x_0^{k-1} + \cdots + a_1x_0)x^{n-1} = (x_0^{k-1} + a_{k-1}x_0^{k-2} + \cdots + a_1)x_0^n$, $A(x_0)$ contains a polynomial of degree $k-1$. By induction method, there exists a polynomial $f(x_0)$ over R^1 such that $f(x_0)x_0^{m+1} = x_0^m$ for some m . Since $x^{n-1} = (-a)p(x)x^{n-1} + x^{n-1} = (-a)(x^{k-1} + a_{k-1}x^{k-2} + \cdots + a_1)x^n + x_0^{n-1}$, we obtain $x^{n+m} = (-a)(x^{k-1} + a_{k-1}x^{k-2} + \cdots + a_1)x^{n+m+1} + x_0^{n+m}$, whence it follows $x^{n+m} = \{(-a)(x^{k-1} + a_{k-1}x^{k-2} + \cdots + a_1) + f(x_0) - ef(x_0)\}x^{n+m+1}$.

Finally, we shall prove the following, which will enable us to readily obtain [24, Theorem 11].

Corollary 10. *Let R^1/R be a left and right integral extension. If R is strongly regular then for any $x \in R^1$ there exists a quasi-regular element u in $R[x]$ such that $x^{2n} - ux^{2n} = x^n$ and $ux^n = x^nu$ for some n .*

Proof. By Theorem 13, there exist s and t in $R[x]$ such that $sx^{n+1} = x^n = x^{n+1}t$ for some n . To be easily seen, $s^n x^{2n} = x^n = x^{2n}t^n$. If we set $a = x^n$ and $b = s^{2n}x^n$, then it is known that $ab = ba$, $a^2b = a$ and $ab^2 = b$ (cf. the proof of [1, Lemma 1]). Obviously, $e = ab$ is an idempotent and $u = e - b$ is a quasi-regular element in $R[x]$ with quasi-inverse $e - a$. Now, it is easy to see that $a^2 - ua^2 = a$.

REFERENCES

- [1] G. AZUMAYA: Strongly π -regular rings, J. Fac. Sci. Hokkaido Univ. Ser. I, **13** (1954), 34-39.
- [2] G. AZUMAYA: Some properties of TTF -classes, Proc. Conference on Orders, Group Rings and Related Topics, Lecture Notes in Math. **353**, Springer-Verlag, Berlin, 1973, pp. 72-83.
- [3] J. A. BEACHY: A characterization of prime ideals, J. Indian Math. Soc. **37** (1973), 343-345.
- [4] B. BROWN and N. H. MCCOY: The maximal regular ideal of a ring, Proc. Amer. Math.

- Soc. **1** (1950), 165—171.
- [5] K. CHIBA and H. TOMINAGA : On strongly regular rings, Proc. Japan Acad. **49** (1973), 435—437.
- [6] K. CHIBA and H. TOMINAGA : Note on strongly regular rings and P_1 -rings, Proc. Japan Acad, **51** (1975), 259—261.
- [7] R. COURTER : Rings all of whose factor rings are semiprime, Canad. Math. Bull. **12** (1969), 417—426.
- [8] C. FAITH : Lectures on Injective Modules and Quotient Rings, Lecture Notes in Math. **49**, Springer-Verlag, Berlin, 1967.
- [9] J. W. FISHER : Nil subrings with bounded indices of nilpotency, J. Algebra **19** (1971), 509—516.
- [10] J. W. FISHER : Von Neumann regular rings versus V -rings, Ring Theory : Proc. Univ. Oklahoma Conference, Dekker, New York, 1974, pp. 101—119.
- [11] J. W. FISHER and R. L. SNIDER : On the von Neumann regularity of rings with regular prime factor rings, Pacific J. Math. **54** (1974), 135—144.
- [12] V. GUPTA : The maximal right weakly regular ideal of a ring, Glasnik Mat. **9** (1974), 29—33.
- [13] F. HANSEN : Die Existenz der Eins in noetherschen Ringen, Archiv der Math. **25** (1974), 589—590.
- [14] F. HANSEN : On one-sided prime ideals, Pacific J. Math. **58** (1975), 79—85.
- [15] I. N. HERSTEIN : Non-Commutative Rings, Carus Math. Monographs **15**, Amer. Math. Ass., New York, 1968.
- [16] A. V. JATEGAONKAR : Left Principal Ideal Rings, Lecture Notes in Math. **123**, Springer-Verlag, Berlin, 1968.
- [17] I. KAPLANSKY : Fields and Rings, Chicago Lectures in Math., Univ. Chicago, Chicago, 1972.
- [18] K. KISHIMOTO and H. TOMINAGA : On decompositions into simple rings. II, Math. J. Okayama Univ. **18** (1975), 39—41.
- [19] J. LAMBEK : Lectures on Rings and Modules, Blaisdell, Waltham, 1966.
- [20] T. H. LENAGAN : Artinian ideals in Noetherian rings, Proc. Amer. Math. Soc. **51** (1975), 499—500.
- [21] G. MICHLER and O. VILLAMAYOR : On rings whose simple modules are injective, J. Algebra **25** (1973), 185—201.
- [22] V. S. RAMAMURTHI : Weakly regular rings, Canad. Math. Bull. **16** (1973), 317—321.
- [23] H. TOMINAGA : On decompositions into simple rings, Math. J. Okayama Univ. **17** (1975), 159—163.
- [24] E. T. WONG : Almost commutative rings and their integral extensions, Math. J. Okayama Univ. **18** (1976), 105—111.
- [25] R. YUE CHI MING : On simple \mathfrak{p} -injective modules, Math. Japonicae **19** (1974), 173—176.

OKAYAMA UNIVERSITY

(Received November 15, 1975)

Added in proof. Recently, Theorem 3 has been proved also by F. Hansen [Proc. Amer. Math. Soc. 55 (1976), 281—286].