

A NOTE ON COMMUTATIVE SEPARABLE ALGEBRAS. II

DAVID E. DOBBS and STUART SUI-SHENG WANG

In this note, we indicate how to employ results concerning descent of projectivity in order to obtain a new proof of the main result in [3, Theorem], namely, that separability for commutative algebras descends by faithful flatness. Following the proof, we comment on the noncommutative case.

Throughout, rings and algebras have identity elements. As usual, if A is a commutative ring and B is an A -algebra, then B is said to be a separable A -algebra if and only if the multiplication map from $B \otimes_A B^o$ to B induces a projective left $B \otimes_A B^o$ -module structure on B [2, p. 40].

Theorem. *Let B be a commutative A -algebra and C a commutative faithfully flat A -algebra. If the C -algebra $C \otimes_A B$ is separable, then B is separable over A .*

Proof. We set $X = B \otimes_A B^o$, $Y = C \otimes_A (B \otimes_A B^o)$, and $Z = (C \otimes_A B) \otimes_C (C \otimes_A B^o)$. Let p and p' be the multiplication maps $X \rightarrow B$ and $Z \rightarrow C \otimes_A B$ respectively. Moreover, let g and h be the canonical isomorphisms

$$Z \rightarrow Y \quad \text{and} \quad C \otimes_A B \rightarrow Y \otimes_X B$$

respectively. Then, the module $C \otimes_A B$ is a left Z -module (under the p' -structure), and is also a left Y -module (under the $1 \otimes p$ -structure). Since $p' = (1 \otimes p)g$ and h is Y -linear, the left Z -module structure on $C \otimes_A B$ may be identified with the left Y -module structure on $Y \otimes_X B$. By hypothesis, $C \otimes_A B$ is a projective left Z -module; moreover, it is cyclic and, a fortiori, finitely generated. Hence $Y \otimes_X B$ is a finitely generated projective left Y -module. Note that Y is a faithfully flat right X -module since faithful flatness is preserved under change of base ring [1, Ch. I, Prop. 5, p. 31]. Therefore, B is a (finitely generated) projective left X -module by virtue of the descent result [1, Ch. I, Prop. 12, p. 35].

Remark. The proof of the theorem was phrased in a way that suggests an attack on the more general context in which B is assumed to be noncommutative. One then needs only to show that B is a flat left

X -module, given that $Y \otimes_X B$ is a flat (indeed, finitely generated projective) left Y -module and Y is faithfully flat over X (on the left and the right). In this generality the problem remains open.

REFERENCES

- [1] N. BOURBAKI: Commutative Algebra (Translated from the French. Actualités Sci. Ind. 1290, Hermann, Paris), Addison-Wesley (1972). MR 50 # 12997.
- [2] F. DEMEYER and E. INGRAHAM: Separable Algebras over Commutative Rings, Lecture Notes in Math. 181 (1971), Springer. MR 43 # 6199.
- [3] S. S.-S. WANG: A note on commutative separable algebras, Math. J. Okayama Univ. 22 (1980), 1—3. MR 81h: 16013.

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF TENNESSEE
AND
DEPARTMENT OF MATHEMATICAL SCIENCES
OAKLAND UNIVERSITY

(Received September 1, 1980)