

## ON GENERALIZED $n$ -LIKE RINGS AND RELATED RINGS

HISAO TOMINAGA and ADIL YAQUB

Throughout,  $R$  will represent a ring with (Jacobson) radical  $J$ , and  $N$  the set of all nilpotent elements in  $R$ . A ring  $R$  is called an *s-unital ring* if for each  $x \in R$  there holds  $x \in Rx \cap xR$ . If  $R$  is an *s-unital ring* then for any finite subset  $F$  of  $R$  there exists an element  $e$  in  $R$  such that  $ex = xe = x$  for all  $x \in F$  (see, [4, Lemma 1 (a)]). Such an element  $e$  will be called a *pseudo-identity* of  $F$ . A ring  $R$  is called a *generalized  $n$ -like ring* if  $R$  satisfies the polynomial identity  $(xy)^n - xy^n - x^n y + xy = 0$  for an integer  $n > 1$ . Recently, H. G. Moore [3] showed that if  $n$  is even or 3 then every generalized  $n$ -like ring with identity is commutative.

The present objective is to prove a theorem which generalizes Theorem 4 of [3] and deduces Theorems 2 and 3 of [3]. We begin with the following lemmas.

**Lemma 1.** *Suppose that for each pair of elements  $x, y$  in  $R$  there exists an integer  $n = n(x, y) > 1$  such that*

$$(*) \quad (xy)^n - xy^n - x^n y + xy = 0.$$

*Then there holds the following:*

- (1)  $(x^{n(x,x)} - x)^2 = x^{2n(x,x)} - 2x^{n(x,x)+1} + x^2 = 0$ .
- (2)  $x^{k(n(x,x)-1)+2} = k(x^{n(x,x)+1} - x^2) + x^2$  for all positive integers  $k$ .
- (3) *If  $R$  is semi-primitive then  $R$  is commutative.*
- (4)  $N^2 = 0$  and  $N = J$  contains the commutator ideal of  $R$ .

*Proof.* (1) Setting  $y = x$  in (\*), we get (1).

(2) Let  $m = n(x, x)$ . Suppose  $x^{k(m-1)+2} = kx^{m+1} - (k-1)x^2$ . Then, by (1),

$$\begin{aligned} x^{(k+1)(m-1)+2} &= x^{m-1} x^{k(m-1)+2} = kx^{2m} - (k-1)x^{m+1} \\ &= k(2x^{m+1} - x^2) - (k-1)x^{m+1} = (k+1)x^{m+1} - kx^2, \end{aligned}$$

which completes the induction.

(3) Note that our hypothesis is inherited by all subrings and homomorphic images of  $R$ . Note also that no complete matrix ring  $(S)_t$  over a division ring  $S$  ( $t > 1$ ) satisfies the hypothesis, as a consideration of  $x = E_{11} + E_{12}$  and  $y = E_{22}$  shows. Because of these facts and the structure

theory of primitive rings, we may assume that  $R$  is a division ring. Then, since  $x^{n(x,x)} - x = 0$  by (1), a well-known theorem of Jacobson shows that  $R$  is commutative.

(4) Since  $x^2 = x^2(2x^{n(x,x)-1} - x^{2(n(x,x)-1)})$  by (1), we see that  $J$  is a nil ideal and every nilpotent element of  $R$  squares to 0. By (3),  $R/J$  is commutative. Hence  $J$  coincides with  $N$  and contains the commutator ideal of  $R$ . Finally, if  $u, v$  are in  $J$  then  $uv = uv^{n(u,v)} + u^{n(u,v)}v - (uv)^{n(u,v)} = 0$ .

**Lemma 2.** *Let  $R$  be an  $s$ -unital ring satisfying the hypothesis in Lemma 1. Then there holds the following:*

(1) *For each  $x \in R$  there exists a positive integer  $\alpha$  such that  $x^{\alpha(n(x,x)-1)}$  is an idempotent.*

(2) *Every idempotent of  $R$  is central.*

*Proof.* (1) Let  $e$  be a pseudo-identity of  $x$ , and set  $\alpha = (2^{n(2e,2e)} - 2)^2$ . Then, by Lemma 1 (1), we get  $0 = ((2e)^{n(2e,2e)} - 2e)^2x = \alpha x$ . Thus, Lemma 1 (2) shows that  $x^{\alpha(n(x,x)-1)+2} = x^2$ , whence (1) follows.

(2) Let  $a, b$  be idempotents in  $R$ , and  $e$  a pseudo-identity of  $\{a, b\}$ . According to (1), we may assume that  $e$  itself is an idempotent. We set  $l = n((e-a)b, a)$  and  $m = n(e-a, b)$ . Then, by (\*),

$$\{(e-a)b\}^l a = \{(e-a)ba\}^l - (e-a)ba^l + (e-a)ba = 0.$$

But, again by (\*),

$$\{(e-a)b\}^m = (e-a)b^m + (e-a)^m b - (e-a)b = (e-a)b,$$

and therefore  $\{(e-a)b\}^m a = (e-a)ba$ . Reiterating in the last and using  $\{(e-a)b\}^l a = 0$  above, we get  $(e-a)ba = 0$ , and hence  $ba = aba$ . Replacing  $a$  by the idempotent  $e-a$  in the above argument, we also have  $b(e-a) = (e-a)b(e-a)$ , and hence  $ab = aba$ . Combining these, we conclude that  $ab = ba$ , and thus all idempotents of  $R$  are central.

**Lemma 3.** (1)  *$R$  is a generalized  $n$ -like ring if and only if  $R$  satisfies the polynomial identities  $(xy)^n = x^n y^n$  and  $(x^n - x)(y^n - y) = 0$ .*

(2) *If  $R$  is an  $s$ -unital generalized  $n$ -like ring then  $(n-1)[u, x] = 0$  for all  $u \in N$  and  $x \in R$ .*

*Proof.* (1) If  $R$  is a generalized  $n$ -like ring, then  $R$  satisfies the polynomial identity  $x^n y^n - xy^n - x^n y + xy = (x^n - x)(y^n - y) = 0$  (Lemma 1 (1) and (4)). Combining this with  $(xy)^n - xy^n - x^n y + xy = 0$ , we readily obtain  $(xy)^n = x^n y^n$ . The converse is trivial.

(2) According to Lemma 1 (4), we have

$$0 = \{(xu)^n - xu^n - x^n u + xu\} - \{(ux)^n - ux^n - u^n x + ux\} = [u, x^n] - [u, x].$$

Now, let  $e$  be a pseudo-identity of  $\{x, u\}$ . Then, by (1) and Lemma 1 (4),

$$\begin{aligned} [u, x] &= [u, x^n] = (ux + x)^n - (xu + x)^n = \{(u + e)x\}^n - \{x(u + e)\}^n \\ &= [(u + e)^n, x^n] = n[u, x^n] = n[u, x], \end{aligned}$$

which implies (2).

We are now in a position to state our main theorem.

**Theorem 1.** *Let  $R$  be an  $s$ -unital (directly) indecomposable ring. Suppose that for each pair of elements  $x, y$  in  $R$  there exists an integer  $n = n(x, y) > 1$  such that  $(xy)^n - xy^n - x^n y + xy = 0$ . Then  $R$  is a local ring whose characteristic is  $p$  or  $p^2$ ,  $p$  a prime.*

*Proof.* Since  $R$  is indecomposable, Lemma 1 (4) and Lemma 2 show that  $R$  contains 1 and is a local ring. Moreover, noting that  $(2^{n(2,2)} - 2)^2 = 0$  by Lemma 1 (1), we see that the characteristic of  $R$  is a power of a prime  $p$ . Since  $p$  is in  $N$ , we get  $p^2 = 0$  (Lemma 1 (4)).

**Corollary 1.** *Let  $R$  be an  $s$ -unital ring. Suppose that for each pair of elements  $x, y$  in  $R$  there exists an integer  $n = n(x, y) > 1$  such that  $(xy)^n - xy^n - x^n y + xy = 0$ . Then  $R$  is a subdirect sum of local rings. If furthermore  $[xy, yx] = 0$  for all  $x \in N$  and  $y \in N$ , then  $R$  is commutative.*

*Proof.* In view of Theorem 1, it remains only to prove the latter part. Note that if  $R^*$  is a homomorphic image of  $R$  then  $[x^* y^*, y^* x^*] = 0$  for all non-nilpotent elements  $x^*, y^*$  in  $R^*$ . Because of this fact, we may assume that  $R$  is subdirectly irreducible, and thus  $R$  is a local ring (Theorem 1). Then, noting that  $N$  is commutative (Lemma 1 (4)), we can easily see that  $[xy, yx] = 0$  for all  $x, y \in R$ . Hence,

$$[x, [x, y]] = [x(y + 1), [x, y + 1]] - [xy, [x, y]] = 0.$$

Now, by [2, Theorem], we see that  $R$  is commutative.

**Corollary 2.** *Let  $R$  be an  $s$ -unital generalized  $n$ -like ring. If  $R$  is indecomposable then  $R$  is a local ring whose characteristic is  $p$  or  $p^2$ ,  $p$  a prime; if  $p$  does not divide  $n - 1$  then  $R$  is commutative.*

*Proof.* In view of Theorem 1, it remains only to prove that if  $(p, n - 1) = 1$  then  $R$  is commutative. By Lemma 3 (2),  $(n - 1)[u, x] = 0$  for all

$u \in N$  and  $x \in R$ . Combining this with  $p^2[u, x] = 0$ , we obtain  $[u, x] = 0$ , and thus  $N$  is contained in the center of  $R$ . Then, using Lemma 1 (1) and [1, Theorem], we see that  $R$  is commutative.

The next includes Theorems 2 and 3 of [3].

**Corollary 3.** *Let  $R$  be an  $s$ -unital generalized  $n$ -like ring. If  $n$  is even or 3, then  $R$  is commutative.*

*Proof.* Without loss of generality, we may assume that  $R$  is subdirectly irreducible, and therefore  $R$  is a local ring by Theorem 1. If  $n$  is even, then  $4 = \{(-1)^n - (-1)\}^2 = 0$  (Lemma 1 (1)). Hence  $R$  is commutative by Corollary 2. Next, we consider the case that  $n = 3$ . Since  $R$  is a local ring, it is enough to show that if  $x, y$  are units in  $R$  then  $xy = yx$ . By Lemma 3 (1),

$$x^2y^2 - x^2 - y^2 + 1 = x^{-1}(x^3 - x)(y^3 - y)y^{-1} = 0$$

and  $y^2x^2 - y^2 - x^2 + 1 = 0$ . Hence  $x^2y^2 = y^2x^2$ . Using this and Lemma 3 (1), we get

$$(xy)^3 = x^3y^3 = xx^2y^2y = xy^2x^2y = (xy)(yx)(xy),$$

whence it follows that  $xy = yx$ .

**Remark.** H. G. Moore required a theorem of Herstein [1] in the proof of [3, Theorem 3]. However, we can prove the same without making use of Herstein theorem (see the proof of Corollary 3).

#### REFERENCES

- [1] I. N. HERSTEIN: A generalization of a theorem of Jacobson, Amer. J. Math. 75 (1953), 105–111.
- [2] S. IKEHATA and H. TOMINAGA: A commutativity theorem, Math. Japonica 24 (1979), 29–30.
- [3] H. G. MOORE: Generalized  $n$ -like rings and commutativity, Canad. Math. Bull. 23 (1980), 449–452.
- [4] I. MOGAMI and M. HONGAN: Note on commutativity of rings, Math. J. Okayama Univ. 20 (1978), 21–24.

OKAYAMA UNIVERSITY, OKAYAMA, JAPAN  
UNIVERSITY OF CALIFORNIA, SANTA BARBARA, U.S.A.

(Received April 27, 1981)