

LIE IDEALS AND SEMI-DERIVATIONS OF PRIME RINGS

Dedicated to Professor Hisao Tominaga on his 60th birthday

MOTOSHI HONGAN

Throughout, R will represent a prime ring of characteristic not 2 with center C . Let $d : r \mapsto r'$ be a *non-zero* semi-derivation of R associated with a surjective function $\sigma : R \rightarrow R$, namely d be a non-zero additive endomorphism of R such that $d\sigma = \sigma d$ and $(xy)' = x'\sigma(y) + xy' = x'y + \sigma(x)y$ for all $x, y \in R$. Then, as was shown in [3, Theorem 1], σ becomes a ring homomorphism. Let U be a Lie ideal of R , and put $W = [U, U]$ and $S = [W, W]$. Obviously, $\sigma(U)$ and both W and S are Lie ideals of R . We consider the following conditions :

- (1- U) $U \subseteq C$.
- (2- U) $U' \subseteq C$.
- (3- U) $U'' \subseteq C$.
- (4- U) $[U', U'] \subseteq C$.
- (5- U) There exists some $a \in R \setminus C$ such that $[a, U'] \subseteq C$.
- (6- U) There exists a non-zero $a \in R$ such that $aU' \subseteq C$.
- (7- U) $[u, u'] \in C$ for all $u \in U$.
- (8- U) There exists a non-zero semi-derivation $\delta : r \mapsto r^*$ of R associated with a bijective function $\tau : R \rightarrow R$ such that $\delta\sigma = \sigma\delta$ and $(U')^* \subseteq C$.

All the results in [1] and [5] proved for Lie ideals of prime rings of characteristic not 2 with derivations have been summarized in Theorem A of [4]. The objective of this paper is to prove the following theorem, which will lead us to a generalization of [4, Theorem A] (Corollary 1).

Theorem 1. *(1- U), (1- W), (1- S) and (2- U) are equivalent. and each of (3- U) – (8- U) implies (1- $\sigma(U$)).*

In preparation for proving our theorem, we state several preliminary lemmas.

Lemma 1. (1) $W' \subseteq [U, U'] + [\sigma(U), U'] \subseteq U + \sigma(U)$; in particular, if $U' \subseteq C$ then $W' = 0$.

- (2) If $U' = 0$ then $U \subseteq C$.
 (3) $\sigma(a) = a$ for every $a \in R$ with $a' = 0$.

Proof. (1) This is clear by the equality $[u, v]' = [u', v] + [\sigma(u), v']$ ($u, v \in U$).

(2) Since $[U, R'] \subseteq [U, R]' + [U', \sigma(R)] = 0$, [3, Theorem 4] gives $U \subseteq C$.

(3) Let \dagger be an arbitrary element of R . Then $\sigma(a)r' = (ar)'\dagger = ar'\dagger$, and so $(a - \sigma(a))R' = 0$. Hence $a - \sigma(a) = 0$ by [3, Lemma 1].

Lemma 2. $(1-U)$, $(1-W)$, $(1-S)$ and $(2-U)$ are equivalent.

Proof. By [2, Lemma 3], $(1-U)$, $(1-W)$ and $(1-S)$ are equivalent. It is easy to see that $(1-U)$ implies $(2-U)$. If $(2-U)$ is satisfied then $W' = 0$ by Lemma 1 (1). Hence, by Lemma 1 (2), we have $(1-W)$.

Lemma 3. If there exists a non-zero $a \in R$ such that $aU' = 0$ (or $U'a = 0$) then $U \subseteq C$.

Proof. Suppose, to the contrary, that $U \not\subseteq C$. Let $u, v \in U$, and $r \in R$. Then $0 = a[u, v'ru]' = a([u, v'r]u)' = a[u, v'r]'\sigma(u) + a[u, v'r]u' = auv'ru'$, i.e., $auU'ru' = 0$. Since $U' \neq 0$ (Lemma 1 (2)), by making use of Brauer's trick, we can easily see that $aUU' = 0$. But this is impossible by [2, Lemma 4].

Lemma 4. If $U'' = 0$ then $U \subseteq C$.

Proof. If not, there exists an ideal M of R such that $[M, R] \subseteq U$ but $[M, R] \not\subseteq C$, by [2, Lemma 1]. Let $m \in [M, R]$, $w \in W$ and $r \in R$. Note that $m, [m, r] \in U$, $\sigma(m') = m'$ (Lemma 1 (3)), $w' \in U'$ and $[w', r] \in U + \sigma(U)$ (Lemma 1 (1)). Then $0 = [mw', r]'' = (m[w', r] + [m, r]w')'' = m''[w', r] + 2\sigma(m')[w', r]' + \sigma^2(m)[w', r]'' + 2\sigma([m, r]')w'' + \sigma^2([m, r])w''' = 2m'[w', r]'$, i.e., $0 = [M, R]'[W', R]' = [M, R]'[W', R']$. Since $[M, R] \not\subseteq C$, Lemma 3 gives $[W', R'] = 0$, and so $W' \subseteq C$ by [3, Theorem 4]. But this contradicts Lemma 2.

Lemma 5. (1) If $C' \neq 0$ and $[a, U'] \subseteq C$ for some $a \in R \setminus C$, then $U \subseteq C$.

(2) If $[a, U'] = 0$ for some $a \in R \setminus C$, then $\sigma(U) \subseteq C$.

Proof. (1) There exists a $c \in C$ such that $c' \neq 0$. Then, for any

$u \in U$ and $r \in R$, $c'[a, [u, r]] = [a, [u, cr]'] - \sigma(c)[a, [u, r]'] \in C$. Hence $[a, [a, [U, R]]] = 0$, and so [5, Theorem 4] gives $[U, R] \subseteq C$; in particular, $[a, [a, U]] = 0$. Again by [5, Theorem 4], we get $U \subseteq C$.

(2) In view of (1), we may assume that $C' = 0$. Suppose, to the contrary, that $\sigma(U) \not\subseteq C$. Since $0 = [a, [a, u]'] = [a, [a', \sigma(u)]]$ for any $u \in U$, we have $[a, [a', \sigma(U)]] = 0$, and so [5, Theorem 4] proves that $a' \in C$ and $\sigma(a') = a'$ (Lemma 1 (3)). Then, for any $u \in U$, $0 = [a, [a^2, u]'] = [a, [(a^2)', \sigma(u)]] = [a, [a'\sigma(a) + aa', \sigma(u)]] = a'[a, [a + \sigma(a), \sigma(u)]]$, i. e., $a'[a, [a + \sigma(a), \sigma(U)]] = 0$. Hence either $a' = 0$ or $[a, [a + \sigma(a), \sigma(U)]] = 0$. In the latter case, we have $a + \sigma(a) \in C$ by [5, Theorem 4], and $0 = (a + \sigma(a))' = 2a'$. We have thus seen that $a' = 0$, in either case. Now, by [2, Lemma 1], there exists an ideal M of R such that $[M, R] \subseteq U$ but $[M, R] \not\subseteq C$. Let $m \in [M, R] (\subseteq M \cap U)$, and $u \in U$. Then $m[a, [a, u]] = [a, m[a, u]] = [a, m'[a, u] + \sigma(m)([a, u'] + [a', \sigma(u)])] = [a, m'[a, u] + \sigma(m)[a, u]'] = [a, (m[a, u]')] = [a, [ma, u]'] - [a, ([m, u]a)'] = -[a, [m, u]']a = 0$, i. e., $[M, R]'[a, [a, U]] = 0$. Hence, Lemma 3 shows that $[a, [a, U]] = 0$, and therefore $U \subseteq C$ by [5, Theorem 4]. But this is a contradiction.

Lemma 6. *If $\sigma(U) \not\subseteq C$ and $U'' = 0$ then $R'' = 0$.*

Proof. First, we claim that $R'' \subseteq C$. It is easy to see that $S' \subseteq V + \sigma(V)$ and $S'' \subseteq U + \sigma(U) + \sigma^2(U)$ (see Lemma 1 (1)). Then, for any $s \in S$ and $r \in R$, $[s'', r'''] = [s'', r]''' \in [U, r]''' + [\sigma(U), r]''' + [\sigma^2(U), r]''' = 0$ and $[s', r^{(4)}] = 3\sigma([s'', r''']) + [s', r^{(4)}] = 3[s'', \sigma(r''')] + [s', r^{(4)}] = [s', r']''' \in [V, r]''' + [\sigma(V), r]''' = 0$ (see Lemma 1 (3)). Hence $[S', R^{(4)}] = 0$. Since $\sigma(S) \not\subseteq C$ (Lemma 2), Lemma 5 (2) gives $R^{(4)} \subseteq C$. Now, let $u \in U$ and $r \in R$. Then $0 = [u, r]^{(4)} = 6[u'', \sigma^2(r'')] + 4[u', \sigma(r''')] = 3[u'', \sigma^2(r'')] + 3[u', \sigma(r''')]$, so that $0 = 2[u, r']''' - [u, r]^{(4)} = 2[u', \sigma(r''')]$, i. e., $[U', R'''] = 0$. Thus $R'' \subseteq C$ by Lemma 5 (2). In particular, $R''U'' = (RU'')''' \subseteq C$, and so either $R'' = 0$ or $U'' \subseteq C$. If $U'' \subseteq C$ then $r^{(4)}u' = (ru')^{(4)} - 4\sigma(r''')u'' \in C$, i. e., $R^{(4)}U' \subseteq C$. Since $U' \not\subseteq C$ (Lemma 2), we get $R^{(4)} = 0$. Hence $4\sigma(r''')u'' = (ru')^{(4)} - r^{(4)}u' = 0$, i. e., $R''U'' = 0$. Combining this with $U'' \neq 0$ (Lemma 4), we get $R'' = 0$. This completes the proof.

We are now ready to complete the proof of our theorem.

Proof of Theorem 1. By Lemma 2, it remains to prove that each of $(3-U) - (8-U)$ implies $(1 - \sigma(U))$.

$(7-U) \Leftrightarrow (1 - \sigma(U))$. Linearizing the relations $[u, u'] \in C$ and

$[u', \sigma(u)] = [u, u]' - [u, u'] \in C$, we get $[u, v]' - [u', v] \in C$ and $[u', \sigma(v)] - [\sigma(u), v'] \in C$ ($u, v \in U$). Then, by making use of Jacobi identity, we see that $[u, [u', \sigma(v)]] = [u, [\sigma(u), v']] = [u, [u, v]'] - [u, [u', v]] = [u, [u, v]'] - [u', [u, v]] \in C$, i.e., $[u, [u', \sigma(U)]] \subseteq C$. If $\sigma(U) \not\subseteq C$ then $u' \in C$ by [5, Theorem 4]; $U' \subseteq C$. But this contradicts Lemma 2.

(3- U) \Rightarrow (1- $\sigma(U)$). Let $u, v \in U$. Then $2[\sigma(u'), v'] = [u, v]'' \in C$, i.e., $[\sigma(U), U'] \subseteq C$. In case $C' \neq 0$, Lemma 5 (1) proves that R satisfies (1- $\sigma(U)$). Thus, we assume henceforth that $C' = 0$. Suppose, to the contrary, that $\sigma(U) \not\subseteq C$. Since $U''' \subseteq C' = 0$, Lemma 6 proves that $R''' = 0$. Now, let u be an arbitrary element of U with $u'' = 0$. Then, for any $v \in U$, $[u, v]' = [u', v] = [\sigma(u'), v'] \in C$ and $2u'[u', v]' = 2u'[u, v]'' = 2u'\sigma([u, v]') = (u[u, v]')'' = [u, uv]'' \in C$ (Lemma 1 (3)). Hence $[u', U'] = 0$, and $u' \in C$ (Lemma 5 (2)). In what follows, let u be an arbitrary element of U with $u'' \neq 0$. Since $[u, R]'' = 0$, by what we have just shown above we see that $[u', R''] = [u, R'']' \subseteq C$; in particular, $[u', [u', R'']] = 0$. Thus $u''[u', [u', R']] = [u', [u', u''\sigma^2(R')]] \subseteq [u', [u', (uR')'']] + u''[u', [u', \sigma(R'')]] = 0$, whence $[u', [u', R']] = 0$ follows, and therefore $u''[u', [u', R]] = [u', [u', u''R]] \subseteq [u', [u', (u'R)']'] + u''[u', [u', \sigma(R')]] = 0$. Hence $[u', [u', R]] = 0$, and so $u' \in C$ by [5, Theorem 4]. We have thus seen that $U' \subseteq C$, which contradicts Lemma 2.

(5- U) \Rightarrow (1- $\sigma(U)$). Suppose, to the contrary, that $\sigma(U) \not\subseteq C$. Let u, v be arbitrary elements of U . Then $[a, [u'', \sigma(v)]] = [a, [u, v]'] - [a, [u', v']] = [a, [u', v]'] + [u', [v', a]] + [v', [a, u']] \in C$, i.e., $[a, [u'', \sigma(U)]] \subseteq C$. Hence $u'' \in C$ by [5, Theorem 4]; R satisfies (3- U). This is a contradiction.

(4- U) \Rightarrow (1- $\sigma(U)$). (4- U) implies either (2- U) or (5- U), and hence (1- $\sigma(U)$).

(6- U) \Rightarrow (1- $\sigma(U)$). If $a \in C$ then $U \subseteq C$ (Lemma 2). We assume henceforth that $a \notin C$. If there exists a $c \in C$ such that $c' \neq 0$, then $c'a[u, r] = a[u, cr]' - \sigma(c)a[u, r]' \in C$ ($u \in U, r \in R$), and so $a[a, [U, R]] = [a, a[U, R]] = 0$. This implies $[U, R] \subseteq C$ by [2, Lemma 7], and hence $U \subseteq C$ by [2, Lemma 6]. Thus, in what follows, we assume that $C' = 0$. Suppose, to the contrary, that $\sigma(U) \not\subseteq C$. Since $a[a, U'] = [a, aU'] = 0$ and $a[a', \sigma(U)] = a[a, U]' \subseteq C$, we get $a' \in C$ by [5, Theorem 6]. Now, let $M = RWR$. Then $[M, R] \subseteq U$ (see the proof of [2, Lemma 1]). Since $\sigma(W) \not\subseteq C$ (Lemma 2), $\sigma(M) = R\sigma(W)R \neq 0$, and therefore, as is well known, $\sigma(M)$ is not commutative. Furthermore, in view of [2, Lemma 6], we see that $[\sigma(M), R] \not\subseteq C$. Let $m \in [M, R] (\subseteq M \cap U)$. Then $a[a, m]'$

$= 0$ and $a[a, m]a' = a[a, m]' \sigma(a) + a[a, m]a' = a([a, m]a)' = a[a, ma]' \in C$. Hence $(a')^2[a, \sigma(m)] = a'[a, (am)'] = (a'[a, am])' = (a[a, m]a')' = 0$; $(a')^2[a, [\sigma(M), R]] = 0$. Since $[a, [\sigma(M), R]] \not\subseteq C$ by [2, Lemma 6], we get $a' = 0$. Thus $aU'' = (aU')' = 0$ and $aU''' = 0$. Since $\sigma(a) = a$ (Lemma 1 (3)), it is easy to see that $a\sigma(U'') = 0$, $a[U', \sigma(U)]' \subseteq C$ and $a[U', \sigma^2(U)]' \subseteq C$. Noting here that $S'' \subseteq U + \sigma(U) + \sigma^2(U)$ (see Lemma 1 (1)), we can easily see that $aU'S''' = a[U', S'']' \subseteq C$. Since $aU' \neq 0$ (Lemma 3), we get $S''' \subseteq C$, and so $aU'S'' = a[U, S'']' \subseteq C$. Hence R satisfies (3-S). This is impossible by Lemma 2.

$(8-U) \Leftrightarrow (1-\sigma(U))$. Noting that $W' \subseteq U + \sigma(U)$, we can easily see that $[(\sigma(U))^*, W''] = ([U, W']')^* \subseteq C$. If $W'' \subseteq C$ then R satisfies (5- $\sigma(U)$) with respect to δ , and therefore $\tau\sigma(U) \subseteq C$. Hence $\sigma(U) \subseteq C$. On the other hand, if $W'' \subseteq C$ then $\sigma(W) \subseteq C$ by the above, and $\sigma(U) \subseteq C$ (Lemma 2).

Corollary 1. *If R is not commutative and σ is bijective, then each of (2-S)–(8-S) is equivalent to (1-U).*

Remark 1. Obviously, Theorem 1 gives generalizations of Theorems 3–7 and 9 in [3], as well. Furthermore, it is easy to see that if U is either d -stable or σ -stable and there exists a non-zero semi-derivation $\delta: r \mapsto r^*$ of R associated with a surjective function $\tau: R \rightarrow R$ such that $(U')^* \subseteq C$ then $\tau\sigma(U) \subseteq C$. This is a generalization of Theorem 8 in [3].

REFERENCES

- [1] R. AWTAR : Lie structure in prime rings with derivations, Publ. Math. Debrecen 31 (1984), 209–215.
- [2] J. BERGEN, I. N. HERSTEIN and J. W. KERR : Lie ideals and derivations of prime rings, J. Algebra 71 (1981), 259–267.
- [3] J. C. CHANG : On semi-derivations of prime rings, Chinese J. Math. 12 (1984), 255–262.
- [4] Y. HIRANO, H. TOMINAGA and A. TRZEPIZUR : On a theorem of Posner, Math. J. Okayama Univ. 27 (1985), 25–32.
- [5] P. H. LEE and T. K. LEE : Lie ideals of prime rings with derivations, Bull. Inst. Math. Acad. Sinica 11 (1983), 75–80.

TSUYAMA COLLEGE OF TECHNOLOGY

(Received July 1, 1985)