

**TRADING DEGREE FOR DIMENSION
IN THE SECTION CONJECTURE:
THE NON-ABELIAN SHAPIRO LEMMA**

JAKOB STIX

ABSTRACT. This note aims at providing evidence for the section conjecture of anabelian geometry by establishing its behaviour under Weil restriction of scalars. In particular, the étale fundamental group of the Weil restriction is determined by means of a Shapiro Lemma for non-abelian group cohomology.

CONTENTS

1. Introduction and results	29
2. Non-abelian cohomology	32
3. Weil restriction of scalars	41
References	43

1. INTRODUCTION AND RESULTS

Let K be a field with fixed separable closure K^{sep} . The étale fundamental group $\pi_1(X)$ of a geometrically connected variety X/K forms naturally a non-abelian extension of pro-finite groups

$$1 \rightarrow \pi_1(X \otimes K^{\text{sep}}) \rightarrow \pi_1(X) \rightarrow \text{Gal}(K^{\text{sep}}/K) \rightarrow 1,$$

that we abbreviate by $\pi_1(X/K)$.

1.1. The section conjecture. A K -rational point $x \in X(K)$ yields by functoriality a section $\text{Gal}(K^{\text{sep}}/K) \rightarrow \pi_1(X)$ of $\pi_1(X/K)$, with image the decomposition group of a point \tilde{x} above x in the universal pro-étale cover of X . Having neglected base points and due to the choice of \tilde{x} , only the class of a section up to conjugation by elements from $\pi_1(X \otimes K^{\text{sep}})$ is well defined. Let us denote by $S_{\pi_1(X/K)}$ the set of $\pi_1(X \otimes K^{\text{sep}})$ -conjugacy classes of sections of $\pi_1(X/K)$. The section conjecture of Grothendieck's anabelian geometry [Gr83] speculates the following.

Mathematics Subject Classification. 14H30 (Primary) ; 14G05, 20E22 (Secondary) .

Key words and phrases. Section Conjecture, Rational points, Anabelian Geometry, Non-abelian Shapiro Lemma.

The author acknowledges support provided by DFG grant STI576/1-(1+2).

Conjecture 1 (Grothendieck). *The map $X(K) \rightarrow S_{\pi_1(X/K)}$ which sends a rational point to the section given by its conjugacy class of decomposition groups is bijective if K is a number field and X/K is a geometrically connected, smooth, projective curve of genus at least 2.*

There is also a version of the section conjecture for affine curves. Here rational points at infinity will lead to an abundance of additional *cuspidal* sections, see [EH08] and [Sx08b]. But apart from the obvious modification forced upon us by cuspidal sections the conjecture remains the same. The condition on the genus gets replaced by asking the Euler-characteristic to be negative.

A birational version of the section conjecture over p -adic local fields was successfully addressed by Koenigsmann in [Ko05], and later by Pop [Po07] in a truncated version that exploits spectacularly modest pro-finite data to recover rational points.

Only recently evidence for the section conjecture could be found through the first examples of curves which satisfy the conjecture, see [Sx08a] and [HS08], though for the reason of having neither points nor sections. Another source of evidence has been the study of the cycle class of a section as pioneered by [EW08].

The goal of the present paper is to provide evidence for the section conjecture from a different direction.

1.2. Results — trading degree for dimension. The evidence for the section conjecture presented in this note consists in its compliance with Weil restriction of scalars, see Section 3. Of course, in order for this to make sense, we widen the applicability of the conjecture beyond the case of curves.

Let L/K be a finite separable field extension within K^{sep} , so that $\text{Gal}_L = \text{Gal}(K^{\text{sep}}/L)$ is a subgroup of $\text{Gal}_K = \text{Gal}(K^{\text{sep}}/K)$. Let X/L be a quasi-projective, geometrically connected variety and $R_{L/K} X$ its Weil restriction of scalars as a geometrically connected variety over K . In Section 2 we will construct an induction functor for extensions which turns out to describe the fundamental group of the Weil restriction in characteristic 0 as an extension as follows.

Theorem 2 (Theorem 17 in Section 3). *Let K be a field of characteristic 0 or let X/L be projective. Then the fundamental group $\pi_1(R_{L/K} X/K)$ of the Weil restriction $R_{L/K} X$ of scalars is isomorphic to the non-abelian induction $\text{Ind}_{\text{Gal}_L}^{\text{Gal}_K} \pi_1(X/L)$.*

Next, a non-abelian analogue of Shapiro's Lemma yields a description of the set of conjugacy classes of sections for an induction.

Theorem 3 (Corollary 15 in Section 3). *Let $E = [1 \rightarrow N \rightarrow E \rightarrow H \rightarrow 1]$ be an extension with induction $\text{Ind}_H^G(E) = [1 \rightarrow M \rightarrow \text{Ind}_H^G(E) \rightarrow G \rightarrow 1]$ with respect to a subgroup $H \subseteq G$.*

Then N -conjugacy classes of sections of $E \twoheadrightarrow H$ are naturally in bijection with M -conjugacy classes of sections of $\text{Ind}_H^G(E) \twoheadrightarrow G$.

Combining Theorem 2 and Theorem 3 above, we obtain our piece of evidence for the section conjecture. We note in passing, that no special assumption on the geometry of the quasi-projective variety X is used.

Theorem 4. *Let L/K be a finite separable field extension and let X/L be a quasi-projective, geometrically connected variety. Let K have characteristic 0 or let X/L be projective.*

(1) *Let $s \in S_{\pi_1(X/L)}$ and $t \in S_{\pi_1(\mathbb{R}_{L/K} X/K)}$ be sections that correspond to each other under the bijection of Theorem 3. Then s is the section associated to a rational point $x \in X(L)$ if and only if t is the section associated to a rational point $y \in \mathbb{R}_{L/K} X(K)$. In this case we may choose x and y so that they correspond to each other via the identification $\mathbb{R}_{L/K} X(K) = X(L)$.*

(2) *Applying the functor π_1 yields a bijective map $X(L) \rightarrow S_{\pi_1(X/L)}$ if and only if it yields a bijective map $\mathbb{R}_{L/K} X(K) \rightarrow S_{\pi_1(\mathbb{R}_{L/K} X/K)}$.*

Proof: Theorem 2 and Theorem 3 show that there is a natural commutative diagram

$$\begin{array}{ccc} \mathbb{R}_{L/K} X(K) & \xrightarrow{\cong} & X(L) \\ \downarrow & & \downarrow \\ S_{\pi_1(\mathbb{R}_{L/K} X/K)} & \xrightarrow{\cong} & S_{\pi_1(X/L)} \end{array}$$

whose horizontal maps are bijections. □

Corollary 5. *The section conjecture holds for smooth projective curves over number fields if it holds for smooth, projective algebraic $\mathbb{K}(\pi, 1)$ spaces over \mathbb{Q} (see [Sx02] Appendix A), which embed into their Albanese variety and have non-vanishing Euler-Poincaré characteristic.*

This corollary explains the title of the article. We have lowered the degree of the number field in the section conjecture to 1 at the expense of working with varieties of dimension exceeding 1. Of course, these higher dimensional varieties of interest are simply \mathbb{Q} -forms of products of smooth, projective hyperbolic curves and so the trade might be marginal. But on the one hand, we are not required to limit the section conjecture to curves or products of curves, and secondly, the arithmetic of \mathbb{Q} and so presumably also sections over $\text{Gal}_{\mathbb{Q}}$ are arithmetically much simpler than for more general algebraic number fields. For example, the recent modularity results were first proven for representations of the full $\text{Gal}_{\mathbb{Q}}$. So the base has become simpler.

Another argument in favour of the improvement of our situation is the following weak analogue of [Sx08a] Theorem 17.

Proposition 6. *Let X/\mathbb{Q} be a smooth, projective and geometrically connected variety which is an algebraic $K(\pi, 1)$ space such that $\pi_1(X/\mathbb{Q})$ admits a section. Then any Galois invariant line bundle $L \in \text{Pic}_X^0(\mathbb{Q})$ has vanishing Brauer obstruction $b(L) = 0 \in \text{Br}(\mathbb{Q})$, hence belongs to a genuine line bundle on X .*

Proof: The local components of $b(L)$ are obtained by base change to \mathbb{Q}_p or \mathbb{R} which preserves the assumptions. The real section conjecture as in [Sx08a] Theorem 24 applies also for higher dimensional $K(\pi, 1)$ spaces and yields the existence of a real point, and so $b(L)_{\mathbb{R}} = 0$.

Over a p -adic field the argument of [Sx08a] Proposition 12 and Corollary 14 still show that the order of $b(L)_{\mathbb{Q}_p}$ is a power of p .

The result now follows from the global reciprocity which states that the local invariants of $b(L)$ sum up to 0. A sum of summands of prime power order where every prime occurs at most once can only vanish if all the summands vanish. Hence $b(L)$ vanishes by the local global principle for Brauer groups of number fields. \square

2. NON-ABELIAN COHOMOLOGY

2.1. Twisted generalized wreath products. We recall some group theory in order to fix notations and to put it into a form useful for the sequel. Although we will ultimately apply the results in the context of pro-finite groups, we choose to neglect the topology in the presentation, because this frees us from adding an abundance of "continuous" everywhere. However we note, that in the pro-finite case the subgroups in question should be closed subgroups.

2.1.1. Wreath products. The **wreath product** of two groups G and N along a right G -set A is the semidirect product

$$N \wr G := \left(\prod_{\alpha \in A} N \right) \rtimes G$$

with respect to the action of G on $\prod_{\alpha \in A} N$ given by $g \cdot (n_\alpha)_{\alpha \in A} = (n_{\alpha g})_{\alpha \in A}$, see [Hu67] §15.6.

2.1.2. Induction for groups acting on groups. Let G be a group. A **G -group** is a group N together with an action $\vartheta : G \rightarrow \text{Aut}(N)$. For a subgroup H in G , induction is a functor ind_H^G from H -groups to G -groups defined as follows. As a group

$$\text{ind}_H^G(N) = \{f : G \rightarrow N; f(hg) = \vartheta(h)(f(g)) \text{ all } h \in H\}$$

with pointwise multiplication. The G -action on $f \in \text{ind}_H^G(N)$ comes from right translation of the argument, so $(g.f)(\alpha) = f(\alpha g)$.

2.1.3. Twisted generalized wreath products. We recall that the **twisted generalized wreath product** of the group G with subgroup H relative to the H -group N is the semidirect product

$$N \wr_H G := (N, \vartheta) \wr_H G := \text{ind}_H^G(N) \rtimes G$$

with respect to the natural G -action of the induction as a G -group, see [Ne63] §2, [Me95] I.9.3, [Hu67] §15.10 and [Ha99] §1.

2.1.4. Sections and H^1 . Let N be a G -group. A 1-cocycle of G with values in N is a map $a : G \rightarrow N$ such that for all $s, t \in G$ we have

$$a_{st} = a_s(s.a_t).$$

The first non-abelian cohomology $H^1(G, N)$ is the set of equivalence classes of 1-cocycles, where cocycles a and b are equivalent if there is $c \in N$ with $a_s = cb_s(s.c)^{-1}$ for all $s \in G$.

Two sections of $N \rtimes G \rightarrow G$ are equivalent if they differ by conjugation with an element of N . The following lemma is well known and straight forward.

Lemma 7. *The map which sends a 1-cocycle $s \mapsto a_s$ to the section $s \mapsto a_s \cdot s$, where we have identified G with the second factor in $N \rtimes G$ establishes a natural bijection of $H^1(G, N)$ with the set of equivalence classes of sections of $N \rtimes G \rightarrow G$.*

2.1.5. The non-abelian Shapiro Lemma in degree 1. Let N be an H -group for a subgroup $H \subset G$. The restriction of the G -group $\text{ind}_H^G(N)$ to an H -group admits an H -equivariant map

$$\text{ev}_1 : \text{ind}_H^G(N)|_H \rightarrow N$$

by evaluating at 1. The composition of restriction and evaluation at 1 defines the Shapiro map

$$\text{sh}^1 : H^1(G, \text{ind}_H^G(N)) \rightarrow H^1(H, N).$$

Proposition 8. *The Shapiro map $\text{sh}^1 : H^1(G, \text{ind}_H^G(N)) \rightarrow H^1(H, N)$ is bijective.*

Proof: A 1-cocycle $s \mapsto b_s$ for G with values in $\text{ind}_H^G(N)$ is given by $b_{s,t} = b_s(t) \in N$ for all $s, t \in G$ such that (i) $b_{s,ht} = \vartheta(h)(b_{s,t})$ for all $s, t \in G$ and $h \in H$ and (ii) $b_{st,g} = b_{s,g}b_{t,gs}$ for all $s, t, g \in G$. The map sh^1 on the level of cocycles maps b to $h \mapsto b_{h,1}$.

Surjectivity. We choose a set of representatives $Y \subset G$ for $H \backslash G$ and obtain maps $\gamma : G \rightarrow H$ and $y : G \rightarrow Y$ such that $g = \gamma_g y_g$ for all $g \in G$. Let $a : H \rightarrow N$ be a 1-cocycle, in particular $a_1 = 1$. We set

$$b_{s,t} := (a_{\gamma_t} \vartheta(\gamma_t)(a_{y_t}))^{-1} (a_{\gamma_{ts}} \vartheta(\gamma_{ts})(a_{y_{ts}}))$$

and a routine calculation shows that b is a 1-cocycle mapping to a . The cocycle condition (ii) is best checked by noting that our definition of b implies $b_{s,t} = (b_{t,1})^{-1} b_{ts,1}$.

Injectivity. Let b and b' be cocycles with $\text{sh}^1(b) \sim \text{sh}^1(b')$. The function $f(s) = (b'_{s,1})^{-1} b_{s,1}$ for all $s \in G$ is an element of $\text{ind}_H^G(N)$. It follows that $b'_{s,t} = f(t) b_{s,t} f(ts)^{-1}$ for all $s, t \in G$ which translates into $b \sim b'$. \square

Remark 9. (1) An alternative proof is given in [Ho78] Theorem 4 using the interpretation as conjugacy classes of complements in semidirect products.

(2) Proposition 8 speaks about conjugacy classes of sections of a twisted wreath product $N \wr_H G = \text{ind}_H^G(N) \rtimes G$. The assertion appears in [PQ03] Thm 2.6 in the case of wreath products, i.e., trivial action of H on N . The introduction of [PQ03] contains the observation that this is a non-abelian version of Shapiro's Lemma but does not elaborate on this idea further.

2.2. Extensions after Eilenberg and MacLane. We recall the theory of non-abelian extensions of Eilenberg and MacLane from [EMcL47].

2.2.1. Kernel. A **kernel** or more precisely a **G -kernel** is a group N together with an exterior action $\rho : G \rightarrow \text{Out}(N)$ by a group G , where $\text{Out}(N) = \text{Aut}(N)/\text{Inn}(N)$ is the group of exterior automorphisms of N . We denote the set of all G -kernels on N by $\text{K}(G, N) = \text{Hom}(G, \text{Out}(N))$.

2.2.2. Center. Let $\rho : G \rightarrow \text{Out}(N)$ be a G -kernel. The **center** Z of the kernel is the center of N together with its inherited G -action

$$\chi = \chi_\rho : G \rightarrow \text{Aut}(Z).$$

For distinction purposes we may denote by $Z(\chi)$ the G -module Z with module structure given by $\chi : G \rightarrow \text{Aut}(Z)$. The set of kernels ρ with center equal to χ is denoted by $\text{K}(G, N)_\chi$.

2.2.3. Extensions. An **extension** E of a group G by a group N is a short exact sequence

$$1 \rightarrow N \rightarrow E \rightarrow G \rightarrow 1.$$

Isomorphisms of extensions of G by N respect both G and N identically. We denote the set of isomorphism classes of extensions by $\text{Ext}(G, N)$. An

extension E leads to a kernel ρ via the restriction to N of the conjugation by preimages:

$$\begin{array}{ccc} E & \xrightarrow{e(\cdot)e^{-1}|_N} & \text{Aut}(N) \\ \downarrow & & \downarrow \\ G & \xrightarrow{\rho} & \text{Out}(N) \end{array}$$

The set of isomorphy classes of extensions, whose kernel has center χ is denoted by $\text{Ext}(G, N)_\chi$. The map $\text{Ext}(G, N)_\chi \rightarrow \text{K}(G, N)_\chi$ that assigns to each extension its kernel is well defined.

2.2.4. Obstruction theory. A kernel which is the kernel of an extension is called **extendible**. By pullback of

$$1 \rightarrow N/Z \rightarrow \text{Aut}(N) \rightarrow \text{Out}(N) \rightarrow 1$$

a kernel $\rho : G \rightarrow \text{Out}(N)$ determines an extension

$$1 \rightarrow N/Z \rightarrow E_\rho \rightarrow G \rightarrow 1$$

such that a presumptive extension E with kernel ρ by conjugation canonically sits in a diagram

$$\begin{array}{ccccccccc} 1 & \longrightarrow & N & \longrightarrow & E & \longrightarrow & G & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & \parallel & & \\ 1 & \longrightarrow & N/Z & \longrightarrow & E_\rho & \longrightarrow & G & \longrightarrow & 1 \end{array}$$

Let χ be the center of ρ . If N were an abelian group with a G -submodule Z , then the existence of E lifting E_ρ were controlled by the coboundary $\delta(E_\rho)$ under $\delta : \text{H}^2(G, N/Z) \rightarrow \text{H}^3(G, Z(\chi))$. The set of such lifts would receive a transitive action of the group $\text{H}^2(G, Z(\chi))$ via the homomorphism $\text{H}^2(G, Z(\chi)) \rightarrow \text{H}^2(G, N)$. In the non-abelian case discussed here (see [EMcL47] §7+8), as Z is central in N , the same formulas with inhomogeneous cocycles which prove the assertions in the abelian case succeed to give the following result, except for the 0 on both sides which follows from [EMcL47] §9+11.

Proposition 10 (essentially [EMcL47]). *The following is exact*

$$0 \rightarrow \text{H}^2(G, Z(\chi)) \rightarrow \text{Ext}(G, N)_\chi \rightarrow \text{K}(G, N)_\chi \xrightarrow{\delta} \text{H}^3(G, Z(\chi)) \rightarrow 0$$

in the sense that $\text{H}^2(G, Z(\chi))$ acts freely on $\text{Ext}(G, N)_\chi$ with quotient set equal to the set of extendible kernels $\delta^{-1}(0)$ and δ is surjective.

The action of $a \in \text{H}^2(G, Z(\chi))$ on $\text{Ext}(G, N)_\chi$ can be constructed on extensions as follows. Let $1 \rightarrow Z \rightarrow \mathcal{Z}_a \rightarrow G \rightarrow 1$ be an extension realizing the cohomology class $a \in \text{H}^2(G, Z(\chi))$, and let $1 \rightarrow N \rightarrow E \rightarrow G \rightarrow 1$

be an extension with center of its kernel equal to χ . Then $a.E$ equals the isomorphism class of the extension

$$1 \rightarrow N \xrightarrow{\text{inclusion}, 0} (E \times_G \mathcal{L}_a) / \Delta(Z) \rightarrow G \rightarrow 1,$$

where $\Delta : Z \rightarrow N \times Z$ is the antidiagonal $z \mapsto (z, -z)$.

2.2.5. Categories of extensions. Let G be a group. The category $\text{Ext}[G]$ has as objects extensions of G with arbitrary kernel and morphisms are maps of extensions up to composition by inner automorphisms from elements of the kernel.

Pushing an extension $1 \rightarrow N \rightarrow E \rightarrow G \rightarrow 1$ by an automorphism of N determines an action of $\text{Out}(N)$ on $\text{Ext}(G, N)$ such that the set of orbits equals the set $\text{Ext}[G, N]$ of isomorphism classes in the category $\text{Ext}[G]$ of extensions of G by N . The map $\text{Ext}(G, N) \rightarrow \text{K}(G, N)$ becomes $\text{Out}(N)$ -equivariant when $\text{Out}(N)$ acts on $\text{K}(G, N) = \text{Hom}(G, \text{Out}(N))$ by composition with inner automorphisms of $\text{Out}(N)$.

2.3. Wreath product type extensions. In this section we built on the work of Holt [Ho78].

2.3.1. Wreath kernels. Let N be a group and $H \subseteq G$ a subgroup. On

$$M = \text{ind}_H^G(N, 1) = \prod_{\alpha \in H \backslash G} N$$

we have an action of G by

$$g \cdot ((n_\alpha)_{\alpha \in H \backslash G}) = (n_{\alpha g})_{\alpha \in H \backslash G}$$

and an outer action $\text{ind}_H^G(\text{Out}(N), 1) = \prod_{\alpha \in H \backslash G} \text{Out}(N) \rightarrow \text{Out}(M)$ given by

$$(f_\alpha)_{\alpha \in H \backslash G} (n_\alpha)_{\alpha \in H \backslash G} = (f_\alpha(n_\alpha))_{\alpha \in H \backslash G}.$$

The two actions are compatible as follows. For $g \in G$ and

$$(f_\alpha)_\alpha = (f_\alpha)_{\alpha \in H \backslash G} \in \prod_{\alpha \in H \backslash G} \text{Out}(N)$$

we have

$$(g \cdot (f_\alpha)_\alpha \cdot g^{-1})((n_\alpha)_{\alpha \in H \backslash G}) = (f_{\alpha g}(n_\alpha))_{\alpha \in H \backslash G} = (g \cdot (f_\alpha)_\alpha)((n_\alpha)_{\alpha \in H \backslash G})$$

resulting in a homomorphism

$$R : (\text{Out}(N), 1) \wr_H G \rightarrow \text{Out}(M).$$

A **wreath kernel** of G on N is a kernel $\rho : G \rightarrow \text{Out}(M)$ together with a lift along R to a homomorphism $\tilde{\rho} : G \rightarrow (\text{Out}(N), 1) \wr_H G$ which is a section of the projection to G . Such a lift is unique if it exists, as two lifts differ at

most by elements in $\prod_{\alpha \in H \setminus G} \text{Out}(N)$ which injects into $\text{Out}(M)$. The set of wreath kernels is thus a subset $K_{\text{wreath}}(G, H; N) \subseteq K(G, M)$ of the set of all G -kernels on M .

2.3.2. The Center of a wreath kernel. The center of M equals $\text{ind}_H^G(Z, 1) = \prod_{\alpha \in H \setminus G} Z$ where Z is the center of N . Thus the center of a wreath kernel lifts to a homomorphism

$$G \rightarrow (\text{Aut}(Z), 1) \wr_H G,$$

which is a section of the projection to G . The construction in Section 2.1.5 performed on the level of cocycles, namely restriction to H and then evaluation at 1, yields the two surjective maps sh as in the following diagram.

$$(2.1) \quad \begin{array}{ccc} K_{\text{wreath}}(G, H; N) & \xrightarrow{\text{sh}} & K(H, N) \\ \text{center} \downarrow & & \downarrow \text{center} \\ \{\text{Sections of } (\text{Aut}(Z), 1) \wr_H G \rightarrow G\} & \xrightarrow{\text{sh}} & \text{Hom}(H, \text{Aut}(Z)) \end{array}$$

The vertical maps associate to a kernel its center. In particular, the center of M as a G -module under the center of the wreath kernel ρ is nothing but $\text{ind}_H^G(\chi) = \text{ind}_H^G(Z(\chi))$ where χ is the center of the H -kernel $\text{sh}(\rho)$.

2.3.3. Wreath product type extensions. A **wreath product type extension**, see [Ho78] p.464, is an extension of G by $M = \prod_{\alpha \in H \setminus G} N$, the kernel of which is a wreath kernel as above. The set of isomorphism classes of wreath product type extensions is denoted by $\text{Ext}_{\text{wreath}}(G, H; N)$. We denote by $\text{Ext}_{\text{wreath}}(G, H; N)_\chi$ (resp. by $K_{\text{wreath}}(G, H; N)_\chi$) the set of those wreath type extensions whose kernel maps (resp. those wreath kernels which map) under sh to a kernel with center χ .

2.3.4. The Shapiro map for extensions. Let $E = [1 \rightarrow M \rightarrow E \xrightarrow{\text{pr}} G \rightarrow 1]$ be a wreath product type extension. The kernel of the map $\text{ev}_1 : M \rightarrow N$, which evaluates at 1 is a normal subgroup of $E|_H = \text{pr}^{-1}(H)$. We may therefore push the restriction of E to H by the map ev_1 to obtain an extension of H by N , that will be denoted $\text{sh}^2(E)$.

2.3.5. Non-abelian Shapiro Lemma in degree 2.

Proposition 11. *Let H be a subgroup of G . Let N be a group with center Z and H -action χ . We have a commutative ladder with exact columns in*

the sense as in Proposition 10

$$\begin{array}{ccc}
0 & & 0 \\
\downarrow & & \downarrow \\
\mathrm{H}^2(G, \mathrm{ind}_H^G(\chi)) & \xrightarrow[\cong]{\mathrm{sh}^2} & \mathrm{H}^2(H, Z(\chi)) \\
\downarrow & & \downarrow \\
\mathrm{Ext}(G, H; N)_\chi & \xrightarrow{\mathrm{sh}^2} & \mathrm{Ext}(H, N)_\chi \\
\downarrow & & \downarrow \\
\mathrm{K}_{\mathrm{wreath}}(G, H; N)_\chi & \xrightarrow{\mathrm{sh}} & \mathrm{K}(H, N)_\chi \\
\downarrow \delta & & \downarrow \delta \\
\mathrm{H}^3(G, \mathrm{ind}_H^G(\chi)) & \xrightarrow[\cong]{\mathrm{sh}^3} & \mathrm{H}^3(H, Z(\chi)) \\
\downarrow & & \downarrow \\
0 & & 0
\end{array}$$

where the horizontal maps are induced by the respective Shapiro map, are all surjective and the two extremal ones are isomorphisms.

Proof: Exactness of the right column is Proposition 10. The commutativity of the diagram follows by tedious but elementary calculations on cochains. The abelian Shapiro Lemma shows that the two extremal horizontal maps are isomorphisms.

The surjectivity of sh was discussed in (2.1). Exactness of the left column follows again from Proposition 10 besides the surjectivity of δ which follows from a diagram chase. Now the surjectivity of the remaining horizontal map follows again by diagram chase. \square

2.3.6. *Split extensions.* The group $\prod_{\alpha \in H \setminus G} \mathrm{Out}(N)$ acts naturally by pushing on isomorphism classes $\mathrm{Ext}_{\mathrm{wreath}}(G, H; N)$ of wreath product type extensions. Let $\mathrm{Ext}_{\mathrm{wreath}}[G, H; N]$ denote the set of orbits.

Theorem 12 ([Ho78] Thm 3). *The map sh^2 yields a bijection*

$$\mathrm{Ext}_{\mathrm{wreath}}[G, H; N] \xrightarrow{\sim} \mathrm{Ext}[H, N].$$

This theorem by Holt has the following immediate corollary.

Corollary 13. *Let $E \in \mathrm{Ext}_{\mathrm{wreath}}(G, H; N)_\chi$ be a wreath product type extension. Then E splits as an extension of G by $\prod_{\alpha \in H \setminus G} N$ if and only if $\mathrm{sh}^2(E)$ splits as an extension of H by N .*

2.4. Non-abelian induction of extensions.

2.4.1. *Adjoint to restriction.* Let H be a subgroup of G . We are going to construct a right adjoint to the functor restriction from G to H

$$\text{res}_H^G : \text{Ext}[G] \rightarrow \text{Ext}[H], \quad E \mapsto \text{res}_H^G E = E|_H$$

which by analogy is called induction from H to G and denoted by Ind_H^G . By Yoneda, all we have to achieve is a proof of the following theorem.

Theorem 14. *Let $F \in \text{Ext}[H]$ be an extension of H . The functor which sends $T \in \text{Ext}[G]$ to $\text{Hom}_{\text{Ext}[H]}(\text{res}_H^G(T), F)$ is representable.*

A representing object as in the theorem is denoted by $\text{Ind}_H^G(F)$. By definition, adjointness

$$(2.2) \quad \text{Hom}_{\text{Ext}[H]}(\text{res}_H^G(T), F) = \text{Hom}_{\text{Ext}[G]}(T, \text{Ind}_H^G(F))$$

holds naturally, and $F \mapsto \text{Ind}_H^G(F)$ is the sought for non-abelian induction functor.

If we apply adjointness to the extension $1 \rightarrow 1 \rightarrow G \xrightarrow{\text{id}} G \rightarrow 1$ we get the following immediate corollary.

Corollary 15. *Let H be a subgroup of G and let*

$$E = [1 \rightarrow N \rightarrow E \rightarrow H \rightarrow 1]$$

be an extension in $\text{Ext}[H]$ with induction

$$\text{Ind}_H^G(E) = [1 \rightarrow M \rightarrow \text{Ind}_H^G(E) \rightarrow G \rightarrow 1]$$

in $\text{Ext}[G]$.

- (1) *The adjointness map (2.2) describes a bijection between N -conjugacy classes of sections of $E \twoheadrightarrow H$ and M -conjugacy classes of sections of $\text{Ind}_H^G(E) \twoheadrightarrow G$.*
- (2) *In particular, E splits if and only if $\text{Ind}_H^G(E)$ splits.*

2.4.2. *Proof of adjointness.* Let $F \in \text{Ext}[H]$ be an extension of H . By Theorem 12 there exist an extension $E \in \text{Ext}[G]$ which is a wreath product type extension such that $\text{sh}^2(E) \cong F$. We are done if we can find a bijection

$$(2.3) \quad \text{sh} : \text{Hom}_{\text{Ext}[G]}(T, E) \xrightarrow{\sim} \text{Hom}_{\text{Ext}[H]}(\text{res}_H^G(T), \text{sh}^2(E))$$

which is natural in $T = [1 \rightarrow S \rightarrow T \rightarrow G \rightarrow 1]$.

Let $\varphi : T \rightarrow E$ be a morphism of extensions. We define the map $\text{sh}(\varphi)$ as the composition $\text{ev}_1 \circ \text{res}_H^G(\varphi)$ which is the vertical map in the following

diagram.

$$\begin{array}{rcl}
\text{res}_H^G(T) & = & [1 \longrightarrow S \longrightarrow T|_H \longrightarrow H \longrightarrow 1] \\
\downarrow \text{res}_H^G(\varphi) & & \downarrow \quad \downarrow \quad \downarrow \\
\text{res}_H^G(E) & = & [1 \longrightarrow M \longrightarrow E|_H \longrightarrow H \longrightarrow 1] \\
\downarrow \text{ev}_1 & & \downarrow \quad \downarrow \quad \downarrow \\
\text{sh}^2(E) & = & [1 \longrightarrow N \longrightarrow \text{sh}^2(E) \longrightarrow H \longrightarrow 1]
\end{array}$$

We denote the projection $T \rightarrow G$ by pr and its restriction to H by $\text{pr}|_H : T|_H \rightarrow H$. A map $\varphi : T \rightarrow E$ is nothing but an M -conjugacy class of sections of

$$\text{pr}_1^* E = [1 \rightarrow M \rightarrow E \times_G T \rightarrow T \rightarrow 1],$$

which is a wreath product type extension for the subgroup $T|_H \subseteq T$. On the other hand, a map $T|_H \rightarrow \text{sh}^2(E)$ is an N -conjugacy class of sections of

$$(\text{pr}|_H)^*(\text{sh}^2(E)) = [1 \rightarrow N \rightarrow \text{sh}^2(E) \times_H T|_H \rightarrow T|_H \rightarrow 1].$$

Because of the isomorphism $\text{sh}^2(\text{pr}_1^* E) \cong (\text{pr}|_H)^*(\text{sh}^2(E))$, we get that in fact Theorem 14 is equivalent to Corollary 15 with $F = \text{Ind}_H^G(E)$ and $E = \text{sh}^2(F)$, which is what we are going to prove in the sequel. Part (2) has already been shown in Corollary 13. We may therefore assume that the wreath product type extension F splits.

We fix a splitting σ , which allows to lift the kernel $\rho_F : G \rightarrow \text{Out}(M)$ to a homomorphism $\vartheta_F : G \rightarrow \text{Aut}(M)$. By means of ϑ_F the extension F is isomorphic to the semidirect product $M \rtimes_{\vartheta_F} G$. Because the following diagram is a fibre product diagram

$$\begin{array}{ccc}
(\text{Aut}(N), 1) \wr_H G & \longrightarrow & (\text{Out}(N), 1) \wr_H G \\
\downarrow \tilde{R} & & \downarrow R \\
\text{Aut}(M) & \longrightarrow & \text{Out}(M)
\end{array}$$

where the map \tilde{R} is constructed analogously to the map R , the wreath kernel $\tilde{\rho}$, which lifts ρ_F , also lifts to a homomorphism $\tilde{\vartheta} : G \rightarrow (\text{Aut}(N), 1) \wr_H G$.

Restriction of the section σ to H and evaluating at 1 induce the corresponding section of $\text{sh}^2(F)$ and a true action $\vartheta : H \rightarrow \text{Aut}(N)$ by means of which $\text{sh}^2(F) \cong N \rtimes_{\vartheta} H$. But ϑ also equals the evaluation at 1 of the restriction to H of $\tilde{\vartheta}$. Hence M as a G -group via $\tilde{\vartheta}$ is nothing but $\text{ind}_H^G(N)$ for the H -group N via ϑ . The result now follows from Proposition 8. \square

2.5. The anabelian case. Although still a mystery in general, it is widely believed that a group only qualifies to be anabelian when its center is trivial. In this section we will work out under the assumption, that the center Z of N is trivial, how the content of the Sections 2.1 - 2.4 specialises. Proposition 10 yields a bijection

$$\text{Ext}(G, N) \rightarrow \text{K}(G, N)$$

with inverse assigning to a kernel $\rho : G \rightarrow \text{Out}(N)$ the pullback under ρ of the extension $1 \rightarrow N \rightarrow \text{Aut}(N) \rightarrow \text{Out}(N) \rightarrow 1$. Hence all kernels are extendible and each extension is determined by its kernel up to isomorphism.

It follows that an N -conjugacy class of sections of an extension with kernel $\rho : G \rightarrow \text{Out}(N)$ is canonically the same as a lift of ρ to an actual action $\tilde{\rho} : G \rightarrow \text{Aut}(N)$ up to uniform conjugation by elements of N .

With N also $M = \prod_{\alpha \in H \setminus G} N$ has trivial center. The fact that sections of $(\text{Out}(N), 1) \wr_H G \twoheadrightarrow G$ and sections of $\text{Out}(N) \times H \twoheadrightarrow H$ correspond to each other under sh^1 up to conjugation by elements from M (resp. N) thus explains Theorem 12 in this case. The content of Corollary 15 follows by applying the same argument to the semi-direct products

$$(\text{Aut}(N), 1) \wr_H G \twoheadrightarrow G$$

and $\text{Aut}(N) \times H \twoheadrightarrow H$.

3. WEIL RESTRICTION OF SCALARS

We content ourselves with a discussion of Weil restriction of scalars relative a finite separable field extension for quasi-projective varieties.

3.1. Properties of the Weil restriction. Let L/K be a finite separable field extension. Weil restriction of scalars relative L/K is a functor $\mathbf{R}_{L/K}$ from quasi-projective varieties over L to quasi-projective varieties over K that is right adjoint to scalar extension $- \otimes_K L$ and thus defined by an identification

$$\text{Hom}_L(Y \otimes_K L, X) = \text{Hom}_K(Y, \mathbf{R}_{L/K}(X))$$

which is natural for schemes X (resp. Y) quasi-projective over L (resp. K), see [BLR90] VII.6. If X has dimension d then $\mathbf{R}_{L/K}(X)$ has dimension $d[L : K]$.

Let K'/K be another field extension and $L' = L \otimes_K K' = \prod_{\alpha} L'_{\alpha}$ the decomposition of the tensor product in separable K' extensions. Then the following Mackey-formula holds

$$(\mathbf{R}_{L/K}(X)) \otimes_K K' = \prod_{\alpha} \mathbf{R}_{L'_{\alpha}/K'}(X \otimes_L L'_{\alpha}).$$

In particular, with an algebraic closure K^{alg} of K we have

$$(\mathbf{R}_{L/K}(X)) \otimes_K K^{\text{alg}} = \prod_{\sigma} X \otimes_{L,\sigma} K^{\text{alg}},$$

where the product is over all K -embeddings $\sigma : L \rightarrow K^{\text{alg}}$. If K^{alg} contains L a priori, then we have a preferred $\sigma_1 = \text{id}$ which gives a projection

$$\text{pr}_1 : (\mathbf{R}_{L/K}(X)) \otimes_K K^{\text{alg}} \rightarrow X \otimes_L K^{\text{alg}}$$

onto the corresponding factor.

3.2. The fundamental group of the Weil restriction. From now on we work with fields of characteristic 0 or ask the variety X/L to be projective. Under these conditions the fundamental group satisfies a Künneth formula [SGA1] Exp X Cor 1.7, Exp XII Cor 5.2, i.e, the projections yield an isomorphism

$$(3.1) \quad \pi_1(\mathbf{R}_{L/K}(X) \otimes_K K^{\text{alg}}) = \pi_1\left(\prod_{\sigma} X \otimes_{L,\sigma} K^{\text{alg}}\right) \xrightarrow{\sim} \prod_{\sigma} \pi_1(X \otimes_{L,\sigma} K^{\text{alg}}).$$

In the extension $\pi_1(\mathbf{R}_{L/K}(X)/K)$, the outer action by conjugation with lifts of $g \in \text{Gal}_K$ on (3.1) acts as $\pi_1(1 \otimes g^{-1})$ and therefore permutes the factors by mapping $\sigma \in \text{Hom}_K(L, K^{\text{alg}}) = \text{Gal}_K / \text{Gal}_L$ to $g^{-1}\sigma$. Reindexing the product with $\alpha = \sigma^{-1}$ transforms this to the right translation action on $\text{Gal}_L \backslash \text{Gal}_K$. Thus we have established the following proposition.

Proposition 16. *The extension $\pi_1(\mathbf{R}_{L/K}(X)/K)$ is a wreath product type extension with respect to the subgroup $\text{Gal}_L \subseteq \text{Gal}_K$.*

In the identification of the index set of the product with $\text{Gal}_L \backslash \text{Gal}_K$ we have chosen a distinguished embedding of L in K^{alg} . Evaluation at 1 then is nothing but the $\pi_1(\text{pr}_1)$ for

$$\text{pr}_1 : (\mathbf{R}_{L/K}(X)) \otimes_K L \rightarrow X$$

which is the adjoint map for the pair of adjoint functors $\mathbf{R}_{L/K}$ and $- \otimes_K L$ and an L -form of the map pr_1 from above. Consequently we find

$$\text{sh}^2(\pi_1(\mathbf{R}_{L/K}(X)/K)) = \pi_1(X/L)$$

and the following structure theorem for the fundamental group of a Weil restriction of scalars holds.

Theorem 17. *The fundamental group $\pi_1(\mathbf{R}_{L/K} X/K)$ is isomorphic to the non-abelian induction $\text{Ind}_{\text{Gal}_L}^{\text{Gal}_K} \pi_1(X/L)$.*

REFERENCES

- [BLR90] Bosch, S., Lütkebohmert, W., Raynaud, M., *Néron Models*, Ergebnisse der Mathematik und ihre Grenzgebiete **21**, Springer, 1990.
- [EMcL47] Eilenberg, S., MacLane, S., Cohomology theory in abstract groups II. Group extensions with a non-abelian kernel. *Ann. of Math.* **48** (1947), no. 2, 326–341.
- [EH08] Esnault, H., Hai, Ph. H., Packets in Grothendieck’s Section Conjecture, *Advances in Mathematics* **218**, Issue 2, (2008), 395–416.
- [EW08] Esnault, H., Wittenberg, O., Remarks on the pronilpotent completion of the fundamental group, preprint, arXiv:0807.2963v3[math.AG], July/December 2008, to appear in *Moscow Mathematical Journal*.
- [Ha99] Haran, D., Hilbertian fields under separable algebraic extensions, *Inventiones Math.* **137** (1999), 113–126.
- [HS08] Harari, D., Szamuely, T., Galois sections for abelianized fundamental groups, with an Appendix by E. V. Flynn, *Math. Annalen* **344**, No. 4 (2009), 779–800.
- [Ho78] Holt, D. F., Embeddings of group extensions into wreath products, *Quart. J. Math. Oxford Ser. (2)* **29** (1978), no. 116, 463–468.
- [Hu67] Huppert, B., *Endliche Gruppen I*, Grundlehren der Mathematischen Wissenschaften **134**, Springer, 1967, xii+793 pp.
- [Gr83] Grothendieck, A., Brief an Faltings (27/06/1983), in: *Geometric Galois Action 1* (ed. L. Schneps, P. Lochak), LMS Lecture Notes **242**, Cambridge 1997, 49–58.
- [Ko05] Koenigsmann, J., On the ‘section conjecture’ in anabelian geometry, *J. Reine Angew. Math.* **588** (2005), 221–235.
- [Me95] Meldrum, J. D. P., *Wreath products of groups and semigroups*, Pitman Monographs and Surveys in Pure and Applied Mathematics **74**, Longman, Harlow, 1995, xii+324 pp.
- [Ne63] Neumann, B. H., Twisted wreath products of groups, *Arch. Math.* **14** (1963), 1–14.
- [PQ03] Parker, C., Quick, M., Maximal complements in wreath products, *Journal of Algebra* **266** (2003), no.1, 320–337.
- [Po07] Pop, F., On the birational p -adic section conjecture, preprint, August 30, 2007.
- [SGA1] Grothendieck, A., Mme. Raynaud, M., *Revêtements Étales et Groupe Fondamental (SGA 1)*, LNM **224**, Springer, 1971.
- [Sx02] Stix, J., Projective anabelian curves in positive characteristic and descent theory for log étale covers, thesis, *Bonner Mathematische Schriften* **354** (2002).
- [Sx08a] Stix, J., On the period-index problem in light of the section conjecture, preprint, February 2008, arXiv:0802.4125v1[math.AG], to appear in *American Journal of Mathematics*.
- [Sx08b] Stix, J., On cuspidal sections of algebraic fundamental groups, preprint, July 2008, arXiv:math.AG/0809.0017v1.

MATHEMATISCHES INSTITUT, UNIVERSITÄT HEIDELBERG, IM NEUENHEIMER FELD 288,
69120 HEIDELBERG

e-mail address: stix@mathi.uni-heidelberg.de

URL: <http://www.mathi.uni-heidelberg.de/~stix/>

(Received February 16, 2009)

(Revised July 12, 2009)